

Nonlinear Ensemble Transform Filter (NETF) for Ocean Assimilation

J. Tödter (toedter@iau.uni-frankfurt.de)¹, P. Kirchgessner², L. Nerger² and B. Ahrens¹

¹Institute for Atmospheric and Environmental Sciences, Goethe University, Frankfurt/Main, Germany, ²Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

Motivation

In nonlinear systems, the analysis moments of the *local ensemble transform Kalman filter* (LETKF)^[1] are biased due to the Gaussian assumption for prior density and observation. The *particle filter* (PF) performs a non-parametric and Bayesian analysis, but suffers from weight divergence.

Approach: Nonlinear Ensemble Transform Filter (NETF)^[2]

- Creates new, equally-weighted analysis ensemble such that its mean and covariance *exactly* match the Bayesian estimators
- Deterministic square root filter as the ETKF
- Domain localization as in the LETKF
- Outperforms (L)ETKF in Lorenz63/96 tests with small ensembles^[2]

NETF

Analysis ensemble with Bayesian moments^[2] (Monte Carlo estimators)

$$\mathbf{w} = (w^1, \dots, w^m)^T$$

Usual PF weights: $w^i \propto p(\mathbf{y}|\mathbf{x}_f^i)$

$$\mathbf{T} = \sqrt{m} \left(\text{diag}(\mathbf{w}) - \mathbf{w}\mathbf{w}^T \right)^{1/2}$$

→ Identical update mechanism: NETF & ETKF only differ by the explicit entries in \mathbf{T} & \mathbf{w} !

NETF Analysis Step: Analogy to the (L)ETKF

Transform forecast ensemble into analysis ensemble with *exactly* specified mean and covariance:

1. Update mean with **weight vector** \mathbf{w} :


$$\bar{\mathbf{x}}_a = \bar{\mathbf{x}}_f + \mathbf{X}'_f \mathbf{w}$$

2. Update perturbations with **transform matrix** \mathbf{T} :

$$\mathbf{X}'_a = \mathbf{X}'_f \mathbf{T} \mathbf{\Lambda}$$

3. Compose final ensemble:

$$\mathbf{X}_a = \bar{\mathbf{x}}_a + \mathbf{X}'_a$$



ETKF

Analysis ensemble with KF moments (Gaussian assumption)

$$\mathbf{w} = \frac{1}{m-1} \mathbf{T} \mathbf{T}^T \mathbf{Y}'_f{}^T \mathbf{R}^{-1} (\mathbf{y} - \bar{\mathbf{y}}_f)$$


$$\mathbf{T} = \left(\mathbf{I} + \mathbf{Y}'_f{}^T \mathbf{R}^{-1} \mathbf{Y}'_f / (m-1) \right)^{-1/2}$$

Notation
 \mathbf{x} = state vector
 m = ensemble size
 $\mathbf{X}'_{f/a}$ = forecast/analysis ens. matrix = $[\mathbf{x}^1, \dots, \mathbf{x}^m]$
 \mathbf{X} = ens. perturbations
 \mathbf{y} = observation vector
 \mathbf{R} = obs. error covariance
 \mathbf{H} = observation operator
 $\mathbf{Y} = \mathbf{H}\mathbf{X}$, $\bar{\mathbf{y}}$ = mean(\mathbf{Y})
 $p(\mathbf{y}|\mathbf{x})$ = likelihood density
 $\mathbf{\Lambda}$ = random rotation matrix

High-Dimensional Ocean Twin Experiment

Model: NEMO v3.3

- Closed square basin, 0.25°, 5km depth
- Driven by zonal wind
- 74 years spin-up
- DA exp. in year 75



State vector

T, U, V, SSH
(on 121x81x11 grid)

→ **dim(state) ≈ 3.3 · 10⁵**


Dynamics

- Double gyre circulation
- Central jet
- Mesoscale eddies
- as e.g. in North Atlantic

→ Realistic & challenging assimilation experiment

Artificial observations^[3]

- each 2nd day → 180 analysis steps
- SSH on *Envisat* tracks
- *Argo* temperature profiles on 3°x3° grid
- **dim(obs) ≈ 3300**




Filter setup

- Localization radius: 2.5° (on average 100 observations per ocean column)
- Inflation factor: 1.025

Initial ensemble

- **dim(ens) = 120**
- from model climatology
- no information about true flow at t=0




→ Generic NETF, no model-dependent changes

Results and Evaluation^[4]

Qualitative evaluation

Snapshots of SSH [m] on day 260:




→ NETF reproduces the true circulation

Quantitative evaluation

RMSEs (normalized at t=0):

- strong error reduction with time compared to free run
- holds for observed (T, SSH) and hidden variables (U, V)
- filter remains stable

→ NETF successfully assimilates the observations




Relative RMSEs for all variables for NETF (full) and free run (dashed)

Comparison to LETKF

with CRPS (averaged over T,U,V,SSH)
Considers *entire ensemble distributions*

- NETF requires a longer spin-up phase than LETKF
- But: better score after convergence




→ Potential benefits of nonlinear analysis

Conclusions and Outlook

Conclusions

- Promising nonlinear filter for high-dim. assimilation
- Simple implementation: analog to (L)ETKF
- Works well in Lorenz to ocean models with small ensemble sizes: *overcomes curse of dimensionality*

Successful development



Future work

- ▶ More large-scale applications
- ▶ Comparison to EWPF
- ▶ Extension to nonlin. smoother
- ▶ ...

References

- [1] Hunt, B. R., E. Kostelich, I. Szunyogh (2007): Efficient data assimilation for spatiotemporal chaos: A local ensemble transform Kalman filter. *Physica D*, 230, 112-126.
- [2] Tödter, J., B. Ahrens (2015): A second-order exact ensemble square root filter for nonlinear data assimilation. *MWR*, 143, 1347-1367.
- [3] Yan, Y., A. Barth, J. M. Beckers (2014): Comparison of different assimilation schemes in a sequential Kalman filter assimilation system. *Ocean Modelling*, 73, 123-137.
- [4] Tödter, J., P. Kirchgessner, L. Nerger, B. Ahrens (2015): Assessment of a nonlinear ensemble transform filter for high-dimensional data assimilation. *MWR*, under review.

Presented at the *Marine Environmental Monitoring, Modelling And Prediction Symposium* (Liège, May 2015)

Support by the projects MiKlip (BMBF, Germany) and SANGOMA (EU) is acknowledged.