

PAPER • OPEN ACCESS

## Global distribution of the cold-water coral *Lophelia pertusa*

To cite this article: R Tong *et al* 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **1004** 012010

View the [article online](#) for updates and enhancements.

You may also like

- [Cold water swelling starch: methods to prepare and recent applications](#)  
Achmat Sarifudin, Enny Sholichah, Woro Setiaboma et al.
- [A Migration Model to Predict the Distribution of Scottish Herring and Mackerel](#)  
Minghao Xu, Dongyu Jiang and Zhe Zhang
- [Application of Extended Habitat Suitability Curves for Depth in Nitrica River](#)  
Peter Ivan and Zuzana Stefunkova



The advertisement features a teal background with white and gold text. On the left, the ECS logo is displayed above the text 'The Electrochemical Society Advancing solid state & electrochemical science & technology'. The main text reads '242nd ECS Meeting Oct 9 – 13, 2022 • Atlanta, GA, US Presenting more than 2,400 technical abstracts in 50 symposia'. In the center, there is a portrait of M. Stanley Whittingham next to a gold Nobel Prize medal. To the right, a 'Register now!' button with a checkmark icon is positioned above a photograph of a large audience seated in a lecture hall. Below the audience photo, a smaller image shows a person's hands holding a futuristic, glowing interface with various icons.

**ECS** The Electrochemical Society  
Advancing solid state & electrochemical science & technology

**242nd ECS Meeting**  
Oct 9 – 13, 2022 • Atlanta, GA, US  
Presenting more than 2,400  
technical abstracts in 50 symposia

**ECS Plenary Lecture**  
featuring  
**M. Stanley Whittingham,**  
Binghamton University  
Nobel Laureate –  
2019 Nobel Prize in Chemistry

Register now!

# Global distribution of the cold-water coral *Lophelia pertusa*

R Tong<sup>1\*</sup>, A J Davies<sup>2</sup>, A Purser<sup>3</sup>, X Liu<sup>4</sup> and F Liu<sup>4</sup>

<sup>1</sup> Fujian University of Technology, Fuzhou, China

<sup>2</sup> University of Rhode Island, Kingston, USA

<sup>3</sup> Alfred-Wegener-Institut, Bremerhaven, Germany

<sup>4</sup> Fuzhou University, Fuzhou, China

**Abstract.** *Lophelia pertusa* plays an important role as a major contributor to many cold-water coral reefs, supporting a high diversity of associated benthic and benthopelagic species. Due to the high sensitivity of *L. pertusa* to human activity, it has been classified as indicator species for Vulnerable Marine Ecosystems. However, the global spatial distribution of *L. pertusa* is far from well known. In this study, a database of *L. pertusa* presence data was compiled derived from the large number of *L. pertusa* occurrence records appearing in recent years. In conjunction with data layers covering a range of environmental drivers, habitat suitability for *L. pertusa* was predicted using the Random Forest approach. Suitable habitat for *L. pertusa* was predicted to occur primarily on continental margins, with the most suitable habitat likely to occur in the North East Atlantic and South Eastern United States of America. Aragonite saturation state, temperature and salinity were identified as the most important contributors to the habitat suitability model. Given the high vulnerability of reef-forming cold-water corals to anthropogenic impacts, habitat suitability models are critical in developing worldwide conservation and management strategies for biodiverse and biomass rich cold-water coral ecosystems.

**Key words-** habitat suitability modelling, framework-forming cold-water coral, random forest

## 1. Introduction

Framework-forming cold-water corals play important roles in the ecological functioning of marine ecosystems (Corbera *et al.*, 2019; Matos *et al.*, 2021). They construct complicated 3-dimensional skeletons of calcium carbonate that may form mounds or reefs, which can fulfil the role of providing an appropriately hard substrate for the settlement of larvae of sessile animals. Cold-water coral reefs also serve as feeding sites, breeding and nursery habitats, and refuges from predators for benthic species, significantly enhancing local biomass and biodiversity (Capezzuto *et al.*, 2018; Freiwald and Roberts, 2005; Linley *et al.*, 2017; Buhl-Mortensen, 2017; Buhl-Mortensen *et al.*, 2017). *Lophelia pertusa* is a scleractinian framework-forming cold-water coral species with a cosmopolitan distribution. It is mainly found at depths from 39 to 3380 m, and is particularly abundant in North Atlantic waters (Roberts *et al.*, 2009; Mortensen, 2000). *Lophelia pertusa* reefs support a high diversity of benthic species, with more than 1300 species found at such reefs in the North East Atlantic (Buhl-Mortensen *et al.*, 2010; Kutti *et al.*, 2014).

Due to physical fragility and slow recovery rates, reef-forming cold-water corals are very sensitive to anthropogenic disturbance, particularly bottom trawling, and climate change (Pecl *et al.*, 2017; Georgian *et al.*, 2016; Movilla *et al.*, 2014). Spatial distribution information for cold-water corals is necessary for the development of conservation strategies, but at present there remain considerable gaps



in knowledge on the distribution of these species. Habitat suitability modelling is a particularly useful approach for charactering the potential distribution of cold-water corals in deep-sea areas (Georgian *et al.*, 2019; Reiss *et al.*, 2015; Bargain *et al.*, 2018; Burgos *et al.*, 2020; Matos *et al.*, 2021; Tong *et al.*, 2013; Wiltshire *et al.*, 2018; Lecours, 2017), providing important input for the creation of conservation and management plans for deep-sea ecosystems (Matos *et al.*, 2021; Reiss *et al.*, 2015). The global habitat suitability of *L. pertusa* has been predicted in several previous studies; using 2060 records, coarse-resolution 1° environmental data and the Ecological-niche factor analysis (ENFA) method (Davies *et al.*, 2008), and with 863 logged occurrences at 30-arc second resolution environmental data and the Maximum Entropy modelling method (Davies and Guinotte, 2011).

This study aims to create an updated database of occurrence records of the framework-forming cold-water coral *L. pertusa*, and develop new global habitat suitability maps for *L. pertusa*, with the model outputs used to better understand its global potential distribution more accurately than was previously possible with the previously utilized smaller databases.

## 2. Methods

### 2.1. Species data

Georeferenced presence-only records were obtained from peer-reviewed scientific outputs and public databases, including the NOAA Deep Sea Coral Data Portal, the ICES Vulnerable Marine Ecosystems data portal and the Ocean Biogeographic Information System portal (OBIS). In order to reduce potential errors in spatial position of the occurrence records included for subsequent modelling, presence records were processed in four steps. Firstly, for records with neither position accuracy data nor depth information provided were excluded. Secondly, records with position accuracy > 1000 m of linear distance were also excluded. Thirdly, for those records with no position accuracy information, but had depth or depth range data, the provided depth or depth range was compared to that extracted from the gridded bathymetry data, and any record where these depths, or depth range and depth, differed by more than 50 m in absolute depth were excluded. Lastly, the remained records with absolute depth values extracted from bathymetry grids < 40 m were excluded. In order to decrease the influence of sampling bias, only one occurrence record was retained in a single cell (after Davies and Guinotte 2011). In addition, similar numbers of cells were randomly chosen to act as pseudo-absence background data.

### 2.2. Environmental variables

Thirty three environmental variables were applied as candidate predictors in this study, with a cell size of 30 arc-sec in a WGS 84 coordinate system (after Davies and Guinotte 2011). These environmental variables can be summarized into four ecologically relevant groups, including 9 terrain variables and gridded bathymetry, 14 water chemistry variables, 7 biological variables, and 2 hydrographic variables (Table 2).

Five environmental variables, namely depth, bottom temperature, aragonite saturation state, bottom horizontal current velocity and vertical current velocity were a priori selected for inclusion in the model, due to their proven high ecological relevance in previous habitat suitability studies for framework-forming cold-water corals published in recent years (Matos *et al.*, 2021; Davies and Guinotte, 2011; Sundahl *et al.*, 2020; Bargain *et al.*, 2018; Dolan *et al.*, 2021; Georgian *et al.*, 2019; Rengstorf *et al.*, 2012).

Since inclusion of highly correlated environmental variables may inhibit model performance and interpretation (Huang *et al.*, 2011), the correlation of environmental variables was investigated by each variable group using analysis of the variance inflation factor (VIF) approach, with depth, temperature, aragonite saturation state, bottom horizontal current velocity and vertical current velocity excluded from the VIF calculations. In addition to the 5 environmental variables, these variables with VIF < 5 indicating a low level of co-linearity, were included in final model (Yesson *et al.*, 2017).

### 2.3. Predictive modeling approach

In this study, the Random Forest for classification and regression approach (RF) was used to predict the global habitat suitability for *L. pertusa*. RF is a regression algorithm, which builds a number of regression trees, and further averages these trees (Breiman, 2001). These trees grow based on different training data subsets created by the bagging procedure to avoid the correlation of trees (Rodriguez-Galiano *et al.*, 2015). RF has been applied to predict benthic species distributions in a series of studies, demonstrating good predictive performances (Matos *et al.*, 2021; Morato *et al.*, 2020; Georgian *et al.*, 2019; Rowden *et al.*, 2017; Gullage *et al.*, 2017).


The species presence and pseudo-absence dataset was randomly split into two sections, 80% as calibration data, with the remaining 20% as evaluation data. Three widely used statistics methods, TSS (true skill statistics), AUC (area under the receiver operating characteristic curve) and kappa statistics, were adopted to evaluate model performance (Allouche *et al.*, 2006). To better understand the predicted habitat suitability, three thresholds were applied to reclassify the predicted map, including sensitivity and specificity balanced threshold, a 90% sensitivity threshold and a 95% sensitivity threshold (Thuiller *et al.*, 2003; Tong *et al.*, 2013).

### 3. Results

In total, 31635 records were obtained from papers and public databases (Table 1, Figure 1). Through processing via the four steps to remove inaccurate records, 3660 records were retained for analysis, with duplicate occurrences within a particular cell deriving from two or more input sources removed. In addition, 3660 cells were randomly chosen as pseudo-absence data.

**Table 1.** The number of occurrence records of *L. pertusa* from different resources.

Resources	OBIS	ICES	NOAA	Papers
Occurrence records	16726	433	12654	1822



**Figure 1.** Spatial distribution of the 31635 *L. pertusa* occurrence records used in this study, with data sources illustrated.

**Table 2.** The 33 environmental variables of four groups used in this study after Davies and Guinotte (2011). The VIF values of environmental variables of three groups tested with  $VIF < 5$  in bold. And the variable importance values of the 13 environmental variables used in habitat suitability modelling illustrated.


Groups	Environmental variables	Reference	VIF	Variable importance
Depth & Terrain variables	<b>Depth</b>	Becker <i>et al.</i> (2009)		0.046
	Slope	Jenness (2013)	20.0	
	<b>Aspect</b>	Jenness (2013)	<b>1.0</b>	0.000
	Plan curvature	Jenness (2013)	38.5	
	<b>Profile curvature</b>	Jenness (2013)	<b>2.4</b>	0.000
	Tangential curvature	Jenness (2013)	36.6	
	Roughness	Wilson <i>et al.</i> (2007)	40.3	
	<b>Rugosity</b>	Jenness (2013)	<b>2.2</b>	0.009
	Terrain ruggedness index	Wilson <i>et al.</i> (2007)	42.9	
	<b>Topographic position index</b>	Wilson <i>et al.</i> (2007)	<b>2.7</b>	0.000
Chemistry variables	<b>Temperature</b>	Boyer <i>et al.</i> (2005)		<b>0.088</b>
	<b>Aragonite saturation states</b>	Steinacher <i>et al.</i> (2009)		<b>0.067</b>
	Alkalinity	Steinacher <i>et al.</i> (2009)	37.2	
	Apparent oxygen utilization	Garcia <i>et al.</i> (2006a)	176.9	
	Dissolved inorganic carbon	Steinacher <i>et al.</i> (2009)	54.1	
	Dissolved oxygen	Garcia <i>et al.</i> (2006a)	31.3	
	Nitrate	Garcia <i>et al.</i> (2006b)	43.2	
	Calcite situation states			
	calcite_stein	Steinacher <i>et al.</i> (2009)	18.3	
	calcite_orr	Orr <i>et al.</i> (2005)	11.4	
	<b>Salinity</b>	Boyer <i>et al.</i> (2005)	<b>1.5</b>	0.050
	Percent oxygen saturation	Garcia <i>et al.</i> (2006a)	185.3	
	Phosphate	Garcia <i>et al.</i> (2006b)	35.5	
Silicate	Garcia <i>et al.</i> (2006b)	12.4		
Biological variables	<b>Seasonal variation index</b>	Lutts <i>et al.</i> (2007)	<b>2.4</b>	0.013
	<b>Particulate organic carbon</b>	Lutts <i>et al.</i> (2007)	<b>2.7</b>	0.053
	Vertically generalized productivity model	Behrenfeld and Falkowski (1997)		
	<b>VPGM_min</b>		<b>2.5</b>	0.003
	VPGM_mean		19.6	
	VPGM_max		6.8	
	Primary productivity	NASA Ocean Color		
	PR_min		Inf	
	PR_mean		15.0	
	PR_max		Inf	
Hydrographic variables	<b>Regional current velocity</b>	Carton <i>et al.</i> (2005)		0.007
	<b>Vertical current velocity</b>	Carton <i>et al.</i> (2005)		0.001

Eight out of 28 environmental variables from the three groups were selected with  $VIF < 5$ , indicating low co-linearity, including aspect, profile curvature, rugosity, topographic position index, salinity, seasonal variation index, particulate organic carbon and vertically generalized productivity model VPGM\_min (Table 2).

The model evaluation showed that the model performed well with TSS 0.974, AUC 0.997 and kappa statistics 0.974. Bottom temperature and aragonite saturation state were the most important variables contributing to the RF model (Table 2). RF predicted suitable habitat for *L. pertusa*



throughout the world's oceans (Figure 2a). The suitable habitat was predicted to occur primarily on continental shelves and slopes (Figure 2). The most suitable habitat for *L. pertusa* was predicted to occur in the North East Atlantic and South Eastern America (Figure 2b).



**Figure 2.** (a) The predicted global habitat suitability for *L. pertusa*. (b) The classified habitat suitability map using 3 thresholds, including the sensitivity and specificity balanced threshold, the 90% sensitivity threshold and the 95% sensitivity threshold. (c) the detailed map of the rectangle section of (b) with hillshade.

#### 4. Discussion and Conclusions

The habitat suitability model developed for *L. pertusa* in this study was influenced primarily by temperature and aragonite saturation state, followed by particulate organic carbon and salinity, which is consistent with findings from previous studies into global stony cold-water coral distributions (Davies *et al.*, 2008; Davies and Guinotte, 2011; Tittensor *et al.*, 2009). The most important variable, temperature, is often reported as one of the most important factors for contributing towards a modelled distribution output for cold-water corals (Matos *et al.*, 2021; Yesson *et al.*, 2017; Sundahl *et al.*, 2020; Bargain *et al.*, 2018). Calcium carbonate in form of aragonite is essential for construction of the skeletal framework of reef-forming cold-water corals and strongly limits the spatial distribution of *L. pertusa* (Georgian *et al.*, 2019; Morato *et al.*, 2020). High-resolution current velocity as a variable can significantly enhance the performance of a predictive model, often correlated with cold-water coral distributions (Rengstorf *et al.*, 2012; Dolan *et al.*, 2021; Bargain *et al.*, 2018). Habitat suitability models at global scales for cold-water corals have shown a weak response to bottom currents, particularly when current conditions are averaged over large spatial extents, indicating resolution of the current is critical if the most accurate results are to be achieved (Davies *et al.*, 2008; Davies and Guinotte, 2011; Tittensor *et al.*, 2009; Yesson *et al.*, 2017; Yesson *et al.*, 2012).

In this study, the database of 31635 *L. pertusa* records was constrained by limitations of position accuracy and/or depth data. The increase in published *L. pertusa* occurrence records since 2011 resulted in significantly more records being included in the model runs than has been possible in previous studies (from 863 in Davies and Guinotte (2011) to 31645, almost a 40-fold increase). Global habitat suitability for *L. pertusa* predicted in this study appears greater than found in the previous study by Davies and Guinotte, 2011 (Figure 2a). A number of areas of the global ocean, such as large seamounts, ridges and canyons, were predicted to have higher habitat suitability than in Davies and Guinotte (2011) study (Figure 2a). However, the high suitability of continental shelves and slopes, particularly in the North East Atlantic and South Eastern United States of America, was similar to previous predictions (Davies and Guinotte, 2011) (Figure 2b). Additionally, the high habitat suitability at North East Atlantic and along continental margin was similar to that demonstrated for other cold-water corals, such as Antipatharia, octocorals and other framework-forming corals (Davies *et al.*, 2008; Davies and Guinotte, 2011; Yesson *et al.*, 2017; Yesson *et al.*, 2012). There was a narrow region along the continental shelf of China predicted to have relatively low habitat suitability for *L. pertusa* (Figure 2c).

The spatial resolution of environmental variables is very important for developing useful and accurate habitat suitability models. The global bathymetric dataset (SRTM30; Becker *et al.* 2009) used in this study, had a resolution of 30-arc seconds, approximately 1 km<sup>2</sup> cell resolution. It successfully captures large terrain features such as large canyons and seamounts (>1 km in diameter), whilst missed smaller terrain features such as mounds and ridges < 1 km in diameter (Davies and Guinotte, 2011). However, smaller terrain relief structures are also known to be important areas for deep-sea scleractinian framework-forming corals. The inclusion of the highest resolution bathymetry dataset SRTM15+ available at present (Tozer *et al.*, 2019) should be considered in any future studies, which may largely improve accuracy of habitat suitability modelling and allow for the incorporation of more species records as resolution increases. The predicted habitat suitability in this study may still be influenced by the sampling bias of *L. pertusa* records. Selecting background data reflecting the same bias as the occurrence data may be useful to reduce the effects of sampling bias (Phillips *et al.*, 2009). Using a kernel density estimate of sampling effort to reduce the influence of sampling bias could significantly improve the model performance (Burgos *et al.*, 2020; Georgian *et al.*, 2019). Such methods to reduce the influence of sampling bias should be considered in future to improve the performance of global habitat suitability model predictions.

### Acknowledgments

This research was supported by the National Key R&D Program of China (NO. 2019YFE0127100), National Natural Science Foundation of China (NO. 42006140), Natural Science Foundation of Fujian Province of China (No.2019I0006).

### References

- [1] Allouche O, Tsoar A and Kadmon R 2006 *Journal of Applied Ecology* **43** 1223-32
- [2] Bargain A, Foglini F, Pairaud I, Bonaldo D, Carniel S, Angeletti L, Taviani M, Rochette S and Fabri M C 2018 *Progress in Oceanography* **169** 151-68
- [3] Becker J, Sandwell D, Smith W, Braud J and Binder B 2009 *Marine Geodesy* **32** 355-71
- [4] Behrenfeld M and Falkowski P 1997 *Limnology & Oceanography* **42** 1-20
- [5] Boyer T, Levitus S, Garcia H, Locarnini R A, Stephens C and Antonov J 2005 *International Journal of Climatology* **25** 931-45
- [6] Breiman L 2001 *Machine Learning* **45** 5-32
- [7] Buhl-Mortensen L, Serigstad B, Buhl-Mortensen P, Olsen M N, Ostrowski M, Blazewicz-Paszkowycz M and Appoh E 2017 *Deep-Sea Research Part II-Topical Studies in Oceanography* **137** 148-56
- [8] Buhl-Mortensen L, Vanreusel A, Gooday A, Levin L, Priede I, Buhl-Mortensen P, Gheerardyn H, King N and Raes M 2010 *Mar. Ecol.* **31** 21-50

- [9] Buhl-Mortensen P 2017 *Marine Biology Research* **13** 1027-40
- [10] Burgos J M, Buhl-Mortensen L, Buhl-Mortensen P, Olafsdottir S H, Steingrund P, Ragnarsson S A and Skagseth O 2020 *Frontiers in Marine Science* **7** 1-25
- [11] Capezzuto F, Ancona F, Carlucci R, Carluccio A, Cornacchia L, Maiorano P, Ricci P, Sion L, Tursi A and D'Onghia G 2018 *Rendiconti Lincei. Scienze Fisiche e Naturali* **29**
- [12] Carton J A, Giese B S and Grodsky S A 2005 *Journal of Geophysical Research Oceans* **110**
- [13] Corbera G, Iacono C L, Gracia E, Grinyo J, Pierdomenico M, Huvenne V, Aguilar R and Gili J M 2019 *Progress In Oceanography* **175**
- [14] Davies A, Wisshak, M, Orr, JC, Roberts, JM 2008 *Deep Sea Research Part I: Oceanographic Research Papers* **55** 1048-62
- [15] Davies A J and Guinotte J M 2011 *PLoS ONE* **6** e18483
- [16] Dolan M, Ross R, Albretsen J, Skardhamar J, Gonzalez-Mirelis G, Bellec V, Buhl-Mortensen P and Bjarnadóttir L 2021 *Geosciences* **11** 48
- [17] Freiwald A and Roberts J M 2005 *Cold-Water Corals and Ecosystems*: (Springer)
- [18] Garcia H E, Locarnini R A, Boyer T P and Antonov J I 2006b *World Ocean Atlas 2005* 4
- [19] Garcia H E, R.A. L, T.P. B and J.I. A 2006a *World ocean atlas 2005* 3
- [20] Georgian S E, Anderson O F and Rowden A A 2019 *Fish Res.* **211** 256-74
- [21] Georgian S E, Dupont S, Kurman M, Butler A, Stromberg S M, Larsson A I and Cordes E E 2016 *Marine Ecology* **37** 1345-59
- [22] Gullage L, Devillers R and Edinger E 2017 *Marine Ecology Progress Series* **582** 57-77
- [23] Jenness J 2012 DEM Surface Tools. [http://www.jennessent.com/arcgis/surface\\_area.htm](http://www.jennessent.com/arcgis/surface_area.htm)
- [24] Kutti T, Bergstad O A, Fosså J H and Helle K 2014 *Deep Sea Research Part II: Topical Studies in Oceanography* **99** 122-33
- [25] Lecours V 2017 *Frontiers in Environmental Science* **4** 1-18
- [26] Linley T D, Lavaleye M, Maiorano P, Bergman M, Capezzuto F, Cousins N J, D' Onghia G, Duineveld G, Shields M A, Sion L, Tursi A and Priede I G 2017 *Deep Sea Research Part II: Topical Studies in Oceanography* **145** 8-21
- [27] Lutz M, Caldeira K, Dunbar R and Behrenfeld M 2007 *Journal of Geophysical Research* **112** C10011
- [28] Matos F L, Company J B and Cunha M R 2021 *Deep-Sea Res. Part I-Oceanogr. Res. Pap.* **170** 12
- [29] Morato T, Gonzalez-Irusta J-M, Dominguez-Carrió C, Wei C-L, Davies A, Sweetman A K, Taranto G H, Beazley L, Garcia-Alegre A, Grehan A, Laffargue P, Murillo F J, Sacau M, Vaz S, Kenchington E, Arnaud-Haond S, Callery O, Chimienti G, Cordes E, Egilisdottir H, Freiwald A, Gasbarro R, Gutierrez-Zarate C, Gianni M, Gilkinson K, Wareham Hayes V E, Hebbeln D, Hedges K, Henry L-A, Johnson D, Koen-Alonso M, Lirette C, Mastrototaro F, Menot L, Molodtsova T, Duran Munoz P, Orejas C, Pennino M G, Puerta P, Ragnarsson S a, Ramiro-Sanchez B, Rice J, Rivera J, Roberts J M, Ross S W, Rueda J L, Sampaio I, Snelgrove P, Stirling D, Treble M A, Urrea J, Vad J, van Oevelen D, Watling L, Walkusz W, Wienberg C, Woillez M, Levin L A and Carreiro-Silva M 2020 *Global Change Biology* **26** 2181-2202
- [30] Mortensen P B 2000 *Department of Fisheries and Marine Biology*, (Norway: University of Bergen) pp 1-42
- [31] Movilla J, Orejas C, Calvo E, Gori A, López-Sanz À, Grinyó J, Domínguez-Carrió C and Pelejero C 2014 *Coral Reefs* **33** 675-86
- [32] Orr J C, Fabry V J, Aumont O, Bopp L, Doney S C, Feely R A, Gnanadesikan A, Gruber N, Ishida A, Joos F, Key R M, Lindsay K, Maier-Reimer E, Matear R, Monfray P, Mouchet A, Najjar R G, Plattner G-K, Rodgers K B, Sabine C L, Sarmiento J L, Schlitzer R, Slater R D, Totterdell I J, Weirig M-F, Yamanaka Y and Yool A 2005 *Nature* **437** 681
- [33] Pecl G T, Araújo M B, Bell J D, Blanchard J, Bonebrake T C, Chen I C, Clark T D, Colwell R K, Danielsen F, Evengård B, Falconi L, Ferrier S, Frusher S, Garcia R A, Griffis R B, Hobday A J, Janion-Scheepers C, Jarzyna M A, Jennings S, Lenoir J, Linnetved H I, Martin V Y, McCormack P C, McDonald J, Mitchell N J, Mustonen T, Pandolfi J M, Pettorelli N, Popova E,



- Robinson S A, Scheffers B R, Shaw J D, Sorte C J, Strugnell J M, Sunday J M, Tuanmu M N, Vergés A, Villanueva C, Wernberg T, Wapstra E and Williams S E 2017 *Science* **355**
- [34] Phillips S J, Dudík M, Elith J, Graham C H, Lehmann A, Leathwick J and Ferrier S 2009 *Ecological Applications* **19** 181-97
- [35] Reiss H, Birchenough S, Borja A, Buhl-Mortensen L, Craeymeersch J, Dannheim J, Darr A, Galparsoro I, Gogina M, Neumann H, Populus J, Rengstorf A M, Valle M, Van Hoey G, Zettler M L and Degraer S 2015 *ICES Journal of Marine Science* **72** 297-315
- [36] Rengstorf A M, Mohn C, Brown C, Wisz M S and Grehan A J 2012 *Deep Sea Research Part I: Oceanographic Research Papers* **93** 72-82
- [37] Roberts J M, Wheeler A, Freiwald A and Cairns S 2009 *Cold-Water Corals: The Biology and Geology of Deep-Sea Coral Habitats*. (Cambridge University Press)
- [38] Rodriguez-Galiano V, Sanchez-Castillo M, Chica-Olmo M and Chica-Rivas M 2015 *Ore Geological Reviews* **71** 804-18
- [39] Rowden A A, Anderson O F, Georgian S E, Bowden D A, Clark M R, Pallentin A and Miller A 2017 *Frontiers in Marine Science* **4** 1-19
- [40] Steinacher M, Joos F, Frölicher T L, Plattner G K and Doney S C 2009 *Biogeosciences* **6** 515-33
- [41] Sundahl H, Buhl-Mortensen P and Buhl-Mortensen L 2020 *Frontiers in Marine Science* **7**
- [42] Thuiller W, Vayreda J, Pino J, Sabate S, Lavorel S and Gracia C 2003 *Global Ecology and Biogeography* **12** 313-25
- [43] Tittensor D P, Baco A R, Brewin P E, Clark M R, Consalvey M, Hall-Spencer J, Rowden A A, Schlacher T, Stocks K I and Rogers A D 2009 *Journal of Biogeography* **36** 1111-28
- [44] Tong R, Purser A, Guinan J and Unnithan V 2013 *Ecological Informatics* **13** 123-32
- [45] Tozer B, Sandwell D, Smith W, Olson C, Beale J and Wessel P 2019 *Earth & Space Science* **6** 1847-64
- [46] Wilson M F J, O'Connell B, Brown C, Guinan J C and Grehan A J 2007 *Marine Geodesy* **30** 33
- [47] Wiltshire K H, Tanner J E, Althaus F, Sorokin S J and Williams A 2018 *Deep-Sea Research Part II-Topical Studies in Oceanography* **157** 121-33
- [48] Yesson C, Bedford F, Rogers A D and Taylor M L 2017 *Deep Sea Research Part II: Topical Studies in Oceanography* **145** 79-86
- [49] Yesson C, Taylor M L, Tittensor D P, Davies A J, Guinotte J, Baco A, Black J, Hall-Spencer J M and Rogers A D 2012 *Journal of Biogeography* **39** 1278-92