



OPEN ACCESS

EDITED BY

Ángel Puga-Bernabéu,
University of Granada, Spain

REVIEWED BY

Annette Goetz,
State Authority for Mining, Energy and
Geology, Germany
Wolfram Michael Kürschner,
University of Oslo, Norway

*CORRESPONDENCE

Agostino Merico,
✉ agostino.merico@leibniz-zmt.de

RECEIVED 25 July 2021

ACCEPTED 17 July 2023

PUBLISHED 31 July 2023

CITATION

Hohn S, Dal Corso J, Hoke KL, Thoms S
and Merico A (2023), Environmental and
physiological conditions that led to the
rise of calcifying nannoplankton in the
Late Triassic.

Front. Earth Sci. 11:747059.

doi: 10.3389/feart.2023.747059

COPYRIGHT

© 2023 Hohn, Dal Corso, Hoke, Thoms
and Merico. This is an open-access
article distributed under the terms of the
[Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is
permitted, provided the original author(s)
and the copyright owner(s) are credited
and that the original publication in this
journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Environmental and physiological conditions that led to the rise of calcifying nannoplankton in the Late Triassic

Sönke Hohn¹, Jacopo Dal Corso², Kim Lisa Hoke³, Silke Thoms⁴
and Agostino Merico^{1,5*}

¹Leibniz Centre for Tropical Marine Research (ZMT), Department of Integrated Modelling, Bremen, Germany, ²State Key Laboratory of Biogeology and Environmental Geology, China University of Geosciences, Wuhan, China, ³Department of Biology, Colorado State University, Fort Collins, CO, United States, ⁴Division of Biosciences, Alfred Wegener Institute for Polar and Marine Research (AWI), Bremerhaven, Germany, ⁵School of Science, Constructor University, Bremen, Germany

The rise of calcifying nannoplankton in the Late Triassic (237–201 Ma) had profound consequences for the carbonate buffering capacity of the ocean. Research on the appearance of calcification typically focuses on the reasons that made this process successful on a global scale. The underlying environmental and physiological conditions that led to its appearance, therefore, are still obscure. Here we show that the rise of intracellular calcification is intrinsically linked to cellular calcium metabolism and to the prevention of calcium accumulation in the cytoplasm. Using gene tree analysis, we present evidence that the physiological machinery for calcification was already present in non-calcifying ancestor cells. Additionally, by modelling the energy demands for calcium transport in calcifying and non-calcifying cells, we demonstrate that intracellular calcification does not require additional energy investments. Since all eukaryotic cells export calcium across the plasma membrane, our findings indicate that the onset of intracellular calcification in Earth's history required only the activation of calcium transport proteins during their passage to the plasma membrane. Hence, intracellular calcification could have occurred at any time in the geological past because physiological preconditions were already present, but a combination of favourable environmental factors may have helped calcifying nannoplankton to spread at the beginning of the Mesozoic, a time of catastrophic climate changes and mass extinctions that reshuffled life on Earth.

KEYWORDS

calcareous nannoplankton, unicellular eukaryotes, calcium homeostasis, ion transport, Carnian, coccolithophores

1 Introduction

Pelagic nannoplankton calcifiers appeared in the Late Triassic (Bralower et al., 1991; Bown, 1998; Bown et al., 2004; Onoue and Yoshida, 2010; Gardin et al., 2012; Preto et al., 2013b; a), Figure 1. The first were probably small calcareous dinoflagellates that produced simple calcispheres (on average about 10 μm in diameter) composed of sub-micron scale calcium carbonate crystals, which appeared during the Carnian, about 233 Ma, when modern marine fauna was rising in the aftermath of the most catastrophic mass extinction event of the entire Phanerozoic Eon at the Permian-Triassic boundary

(Janofske, 1992; Onoue and Yoshida, 2010; Preto et al., 2013b; Preto et al., 2013a; Dal Corso et al., 2021), Figure 1. Coccolithophores, which are haptophyte algae and secrete complex ornamented calcite plates (Young, 2003; Young et al., 2005), appeared in the middle-late Norian, about 215–210 Ma (Gardin et al., 2012; Demangel et al., 2020), and spread after the Triassic-Jurassic mass extinction event (e.g., Bown et al., 2004), Figure 1. Before the emergence of pelagic nannoplankton calcification, the biological precipitation of calcium carbonate was dominated by metazoans and microbial mats that calcified extracellularly in a medium isolated from seawater (Lowenstam and Margulis, 1980; Kaźmierczak et al., 1985; Knoll, 2003). In contrast, calcareous nannoplankton, such as calcareous dinoflagellates (calcispheres) and coccolithophores, calcify inside intracellular compartments called vesicles and extrude the accreted crystals via exocytosis (Inouye and Pienaar, 1983; Paasche, 2002; Brownlee et al., 2015; Jantschke et al., 2020). The emergence of intracellular calcification allowed biocalcification to expand into the surface waters of the open ocean, marking the transition from the so-called “Neritan” ocean to the “Cretan” ocean (Zeebe and Westbroek, 2003; Ridgwell, 2005; de Vargas et al., 2007), Figure 1. This expansion produced a much more efficient regulation of the global carbon cycle and changed the marine calcium carbonate saturation state, transforming the entire ocean carbonate chemistry (Zeebe and Westbroek, 2003; Ridgwell, 2005; Ridgwell and Zeebe, 2005). The production of intracellular platelets that are then extruded onto the outer cell surface led to a completely new phenotype with respect to non-calcifying nannoplankton. The emergence of pelagic calcareous nannoplankton is, thus, one of the most important evolutionary innovations in Earth’s history but how and why it originated is still unclear. The mechanisms driving evolutionary innovations are mutation and selection. Therefore, before identifying the benefits that could have selected intracellular calcification to make it a successful innovation at a global scale, it is important to understand the physiological characteristics (or mutations) required for its emergence.

While the evolution of extracellular calcification was likely an adaptive response to rising calcium concentrations in the ocean (Kaźmierczak et al., 1985; Kaźmierczak and Kempe, 2004), the appearance of intracellular calcification does not clearly correlate with changes in ocean chemistry (Figure 1). For example, a temporal coincidence between a Late Triassic drastic drop in atmospheric $p\text{CO}_2$, resulting from global carbon-climate numerical modelling, and the primary radiation of calcareous nannoplankton, inferred from the fossil record, has been observed but a cause-and-effect link between the two events could not be proven (Goddéris et al., 2008).

Moreover, when intracellular calcification emerged, the concentration of calcium in the ocean was approximately 30 mM, dissolved inorganic carbon (DIC) was 5,000 μM , and ocean pH was about 7.8 (Figure 1). These conditions would have led to a carbonate concentration of 160 μM and a calcite saturation state of 10. Although such chemical conditions would be favourable to calcification and could therefore positively select for this innovation, they alone are not sufficient to explain the appearance of intracellular calcification because the ocean experienced similar conditions 80–100 million years before the first evidence of pelagic nannoplankton calcification appears in the geological record (Late

Triassic-Jurassic). The ocean experienced even higher calcium concentrations (from 50 to almost 60 mM) from the Cambrian to the Carboniferous and calcite saturation reached almost 15 in the Carboniferous and Permian. Therefore, it is plausible to assume that the emergence of intracellular calcification was most likely a consequence of random changes in cellular physiology rather than a response to external environmental factors such as changes in ocean carbonate chemistry. Using new energy budget calculations and gene analyses, we investigate here the underlying mechanisms and prerequisites that allowed intracellular calcification to appear. More explicitly, we investigate if the proteins and ion transporters involved in intracellular calcification were present in non-calcifying ancestors or appeared coincidentally with the emergence of intracellular calcification. Furthermore, we explore if intracellular calcification is indeed energetically costly as proposed by Monteiro et al. (2016) so that secondary evolutionary benefits must compensate for the energy invested into calcification. This is done by calculating the amounts of energy required to pump calcium ions into the calcifying vesicles in comparison to maintaining calcium homeostasis in all (calcifying and non-calcifying) cells.

2 Materials and methods

2.1 Analysis of environmental data

We calculated the carbonate chemistry parameters from reconstructed historic ocean pH and DIC (Ridgwell, 2005) using the standard software CO2sys, version 1.01 (Lewis and Wallace, 1998). Calculated ocean carbonate concentrations and reconstructed calcium concentrations (Stanley and Hardie, 1998) were used to determine the calcite saturation state. Atmospheric $p\text{CO}_2$ was obtained from previous reconstructions (Bernier and Kothavala, 2001). We compared the carbonate chemistry parameters with the fossil record of pelagic calcifying nannoplankton (Bown et al., 2004) to identify the environmental conditions under which intracellular calcification emerged (Figure 1). The primary objective of this analysis is to compare, in qualitative terms, the reconstructed carbonate chemistry of the ocean with the appearance of calcareous nannoplankton. A precise, quantitative reconstruction of the carbonate chemistry throughout the Phanerozoic goes beyond the scope of our study. However, from a visual inspection, the data we compiled using literature sources and CO2sys are consistent with recent reconstructions (Witkowski et al., 2018; Montañez et al., 2016).

2.2 Calcium-transporter gene tree analysis

To examine the evolutionary history of calcification-related proteins, we identified proteins across lineages and created gene trees of relevant protein families. We downloaded protein databases from the National Center for Biotechnology Information (NCBI) genome browser for taxa represented in the phylogeny of Read et al. (2013), Supplementary Table S1. To account for differences in annotation completeness in each protein database, we used BLAST to identify homologs of *Gephyrocapsa huxleyi* proteins in each taxon. We searched the *G. huxleyi* protein database for annotated sequences




FIGURE 1

Fossil record of pelagic calcifiers and paleoceanographic carbonate chemistry over the past 550 Myr. The number of marine families and evolutionary faunas throughout the Phanerozoic are from Raup and Sepkoski Jr (1982) and Sepkoski and Ausich (1990). The numbers indicate the 5 major mass extinction events (1 = Ordovician–Silurian Extinction; 2 = Late Devonian Extinction; 3 = Permian–Triassic Extinction; 4 = Triassic–Jurassic Extinction; 5 = Cretaceous–Paleogene Extinction). The number of pelagic nannoplankton calcifiers is reconstructed by Bown et al. (2004). Note, however, that the data in Bown et al. (2004) include pelagic calcifiers in general and not exclusively nannoplankton. The concentration of calcium is taken from Stanley and Hardie (1998) and atmospheric $p\text{CO}_2$ from Berner and Kothavala (2001). Dissolved inorganic carbon (DIC) and ocean pH were reconstructed by Ridgwell (2005). The calcite saturation state (Ω_{calcite}) was calculated, using the standard software CO2sys (Lewis and Wallace, 1998), from Ca^{2+} and CO_3^{2-} based on ocean temperature (Royer et al., 2004). The emergence of intracellular calcification (indicated by the green arrow) is marked by the rise of calcareous dinoflagellates in the early Late Triassic (Preto et al., 2013b) and was followed by the appearance of coccolithophores in the latest Triassic. Unidentified, very rare calcispheres are found in the Silurian, Devonian, and Carboniferous but are of uncertain origin (Kaźmierczak and Kremer, 2005).

in each protein family implicated in calcification (Benner et al., 2013) to use as query sequences (Supplementary Table S2). We then used pBLAST to identify all sequences in the protein database that matched these query sequences with probabilities less than e^{-6} . After discarding duplicates, we aligned the BLAST results for each protein family in Geneious 11.1.3, (Kearse et al., 2012), <http://www.geneious.com>, using the Geneious global alignment with cost matrix Blosum62, gap open penalty 12, and gap extension penalty 3. We created gene trees using Jukes-Cantor genetic distance module and the UPGMA tree building method in Geneious. We then inspected trees for phylogenetic representation and patterns of protein family expansion. Because the BLAST output results for calcium channels included both cation channels and calmodulin, we split these sequences into two subtrees for

better visualization of the relationships among calmodulin-related proteins. Supplementary Figures S1–S7 show the resulting gene trees.

2.3 Calculation of energy invested into intracellular calcification

We calculated the energy required for calcium transport into Golgi-derived vesicles as part of the energy invested to maintain calcium homeostasis. In our calculations, we neglect the costs associated to the transport of golgi vesicles because these vesicles are transported by both calcifying and non-calcifying cells. We also neglect the costs required for the production of polysaccharides

and for exocytosis, the process for the secretion of polysaccharides by haptophytes, because also these costs are incurred by non-calcifying cells and, thus, do not constitute an additional energy requirement for calcifying cells. In order to achieve intracellular calcium homeostasis, total calcium transport out of the cytoplasm has to equal the passive influx via calcium channels. The passive, diffusive uptake via the calcium channels is driven by the difference between the calcium concentration at the cell surface (Ca_{surf}) and the calcium concentration in the cytoplasm (Ca_{cyt}). Thus, knowing the calcium gradient across the plasma membrane, the energy required to achieve calcium homeostasis (E_{ATP}) can be calculated by

$$E_{ATP} = P_{Ca} \cdot A_{cell} \cdot (Ca_{surf} - Ca_{cyt}) \cdot \Psi \frac{ATP}{Ca}, \quad (1)$$

where P_{Ca} is the apparent membrane permeability per cell surface area (A_{cell}) and $\Psi \frac{ATP}{Ca}$ is the amount of ATP required per calcium ion transported. The consumption of one ATP molecule for transporting two calcium ions through the membrane is a functional property of the reaction loop of the Ca-ATPase (Toyoshima and Inesi, 2004; Winther et al., 2013) and is independent if transport is over the vesicle or plasma membrane. Without net calcium influx into the cell, Ca_{surf} is equal to the calcium concentration in the bulk medium (Ca_{out}). However, fluxes of calcium into intracellular compartments, such as vesicles, create a concentration gradient across the cell membrane with $Ca_{surf} < Ca_{out}$. With the total flux of calcium into the vesicles ($Q_{vesicle}$), and thus into the cell, the integration of the spherical diffusion equation in the cell environment gives the following concentration at the cell surface:

$$Ca_{surf} = Ca_{out} - \frac{Q_{vesicle}}{4 \cdot \pi \cdot r_{cell} \cdot D_{Ca}}, \quad (2)$$

where D_{Ca} is the diffusion coefficient of calcium in seawater and r_{cell} is the cell radius. Here, we assume that the transport of calcium into vesicles represents a fixed fraction of total calcium transport (f_v) as compared to the transport over the plasma membrane ($1 - f_v$), i.e.:

$$Q_{vesicle} = f_v \cdot P_{Ca} \cdot A_{cell} \cdot (Ca_{surf} - Ca_{cyt}). \quad (3)$$

In other words, we assume that while calcium is rushing in through the ion channels, ATPases pump calcium back through the cell membrane into the extracellular medium but at the same time also into the vesicles. The volume inside the vesicle, however, can be regarded as outside of the cytoplasm from a thermodynamical point of view. In order to maintain calcium homeostasis, the flux into the cytoplasm has to be balanced by the fluxes out of the cytoplasm. Inserting the expression for $Q_{vesicle}$ into Eq. 2 and subtracting Ca_{cyt} from both sides of Eq. 2 results in an implicit equation for the concentration difference $Ca_{surf} - Ca_{cyt}$, which has the solution:

$$Ca_{surf} - Ca_{cyt} = \frac{Ca_{out} - Ca_{cyt}}{1 + \frac{f_v \cdot P_{Ca} \cdot A_{cell}}{4 \cdot \pi \cdot r_{cell} \cdot D_{Ca}}} \quad (4)$$

Replacing in Eq. 1 the concentration difference $Ca_{surf} - Ca_{cyt}$ with Eq. 4 gives E_{ATP} as a function of the relative fraction of calcium transport into the calcifying vesicles:

$$E_{ATP} = P_{Ca} \cdot A_{cell} \cdot \frac{Ca_{out} - Ca_{cyt}}{1 + \frac{f_v \cdot P_{Ca} \cdot A_{cell}}{4 \cdot \pi \cdot r_{cell} \cdot D_{Ca}}} \cdot \Psi \frac{ATP}{Ca} \quad (5)$$

This equation shows that the higher f_v , the lower the absolute requirements for cellular energy. In order to quantify how much the cellular energy consumption can potentially be reduced by intracellular calcification, we calculate the energy requirements to maintain calcium homeostasis as a function of f_v from 0 (non-calcifying cell) to 1 (Supplementary Figure S8). The coccolithophore *Gephyrocapsa huxleyi* produces approximately 1 coccolith per hour (Paasche, 1962) and one coccolith contains around 22 fmol of $CaCO_3$ (Young and Ziveri, 2000). This results in a calcification flux ($Q_{vesicle}$) of about $6.11 \cdot 10^{-18} \text{ mol s}^{-1}$ (see: Holtz et al., 2013). Since the permeability of the plasma membrane for calcium ions is not known in coccolithophores, we estimate a permeability that is based on the conductance of calcium channels (Lux and Nagy, 1981) and measured densities of calcium channels in cell membranes of snail axons (Krishtal et al., 1981; Tsien, 1983). All the parameter values used to calculate the extra energy required by intracellular calcification are reported in Supplementary Table S3. We provide the numerical code for calculating the energy required to maintain calcium homeostasis by single-celled calcifying nanoplankton as open-source software (<https://github.com/systemsecologygroup/Calcsphere>) so that it can be used, modified, and redistributed freely.

2.4 Calculation of intracellular calcium poisoning

The energy for the Ca^{2+} pumping ATPase is provided by the hydrolysis of ATP at a high ATP/ADP:Pi ratio in the cytosol with a free energy gain, ΔG , of 15 kcal/mol (Meldolesi and Pozzan, 1998). Cooperative binding of 2 Ca^{2+} per ATPase implies a Ca^{2+} /ATP coupling ratio of 2 (Inesi and Tadini-Buoninsegni, 2014). We assume an electroneutral $Ca^{2+}/2H^+$ co-transport for the Ca^{2+} pump and a physiological pH at both sides of the Ca^{2+} pump. If all of the free energy available from ATP hydrolysis is stored in the concentration gradient of Ca^{2+} between the cytosol and its environment, i.e.: $\Delta G = 2RT \cdot \ln(Ca_{surf}/Ca_{cyt})$, the calcium gradient can be up to 10^5 . If the Ca-ATPases work near this thermodynamic limit, the cytosolic Ca^{2+} concentration of $0.1 \mu\text{mol L}^{-1}$ and the gradient of 10^5 yields a Ca^{2+} concentration of 10 mM at the cell surface, which is, in fact, exactly the Ca^{2+} concentration in today's ocean (Langer et al., 2006). Ca^{2+} transport near the thermodynamic limit would have the advantages that: 1) pumping Ca^{2+} from the cytosol to the environment proceeds with a minimum dissipation of energy in ATP consumption, and 2) a steep Ca^{2+} gradient provides fast influx in cell signaling. We therefore postulate that a constant Ca^{2+} -gradient close to the thermodynamic limit represents a part of calcium homeostasis. Moreover, the Ca^{2+} -gradient across the plasma membrane should be quite independent of the bulk Ca^{2+} concentration in seawater.

Assuming that the Ca^{2+} -gradient across the plasma membrane is maintained at 10^5 by the Ca^{2+} -ATPase, the intracellular calcium concentration will respond to variations in extracellular calcium concentration as experienced during Earth's history (Stanley and Hardie, 1998), Figure 1. The intracellular concentration of calcium (Ca_{cyt}), however, generates a constraint on the intracellular phosphate concentration (PO_{cyt}) because calcium and phosphate precipitate to $Ca_3(PO_4)_2$ when the ion product

exceeds the solubility product $K_{sp}^{\text{Ca}_3(\text{PO}_4)_2} = \text{PO}_{\text{cyt}} \cdot \text{Ca}_{\text{cyt}}$. We can therefore calculate the maximum possible intracellular phosphate concentration for any intracellular calcium concentration as follows:

$$\text{PO}_{\text{cyt}} = \sqrt{\frac{K_{sp}^{\text{Ca}_3(\text{PO}_4)_2}}{\text{Ca}_{\text{cyt}}}}. \quad (6)$$

3 Results and discussion

3.1 Calcium metabolism

Calcium ions play important roles in the physiology of eukaryotic cells, including control of DNA synthesis, chromosomal configuration, and signal transduction (Marcum et al., 1978; Boynton et al., 1980; Krishtal et al., 1981; Lux and Nagy, 1981; Tsien, 1983; Poenie et al., 1985; Carafoli, 1987; Steinhardt and Alderton, 1988; Hepler, 1994; Sanders et al., 2002; Verret et al., 2010). Calcium ions enter the cell passively via channels (Gussone et al., 2006; Verret et al., 2010) that allow for a diffusive influx driven by a steep concentration gradient across the plasma membrane (Langer et al., 2006). Excess intracellular Ca^{2+} is then actively removed by calcium-binding proteins (e.g., calmodulin) and transport proteins (Ca^{2+} -ATPase and $\text{Na}^+/\text{Ca}^{2+}$ exchanger) with different calcium affinities and maximum transport rates (Carafoli, 1987). The Ca^{2+} concentration inside the cell has to be tightly regulated (calcium homeostasis) at around $0.1 \mu\text{mol L}^{-1}$ (Brownlee et al., 1995) for optimal cell functioning (Carafoli, 1987) and to avoid cell poisoning (Simkiss, 1977; Orrenius et al., 1989; Müller et al., 2015).

The same proteins that maintain intracellular calcium homeostasis in non-calcifying cells (Figure 2A) are implicated in intracellular calcification (Figure 2B). To determine whether an expansion of protein families coincides with the emergence of intracellular calcification, we compared all gene families involved in the precipitation of calcium carbonate (von Dassow et al., 2009) in a large number of eukaryotes (Read et al., 2013). These genes include Ca^{2+} transporting ATPases, HCO_3^- transporters, Ca^{2+} channels, calmodulin, carbonic anhydrase, $\text{Ca}^{2+}/\text{H}^+$ exchangers, $\text{Na}^+/\text{Ca}^{2+}$ exchangers, and vacuolar H^+ -ATPases. The presence of calcium transport proteins in all eukaryotes clearly indicates that these genes are not unique to intracellular calcification. We found no evidence that diversification of these gene families could explain the appearance of nannoplankton calcification (Supplementary Figures S1–S7).

The most significant difference between calcifying and non-calcifying cells is that calcium carbonate precipitates inside Golgi-derived vesicles. The vesicles fuse with the plasma membrane and the calcium carbonate is extruded outside the cell where it remains attached to the extracellular organic matrix. The same vesicular pathway is used in non-calcifying cells to produce new membrane material and to transport proteins to the plasma membrane (Figure 3). All ion transporters and channels that help establish calcium homeostasis in non-calcifying cells, therefore, reach the plasma membrane via Golgi-derived transport vesicles (Pfeffer and Rothmann, 1987). However, the Ca^{2+} transporting ATPase initially occurs in its auto-inhibited form and becomes activated

by the presence of calmodulin (Tidow et al., 2012) at the plasma membrane. We thus argue that the emergence of intracellular calcification was triggered by the activation of calcium pumps during the passage of transport vesicles from the Golgi apparatus to the plasma membrane, following the accumulation of calcium in the transport vesicles. This explanation is consistent with the simple structure of sub-micron scaled calcite crystals of proto-calcispheres found in previous studies and attributed to calcareous dinoflagellates (Janofske, 1992; Preto et al., 2013b; Dal Corso et al., 2021). The activation of the calcium transport inside the Golgi apparatus would later lead to the formation of more elaborated coccolith-like platelets due to the specific endomembranous structure of the Golgi apparatus (Figure 3). This is also in line with the finding that inhibition of the cytoskeleton and microtubule formation induces malformations in modern coccolith (Langer et al., 2010).

Such a change in timing of activation would not be detected in gene sequence comparison, as they require protein functional assays. We conclude, however, that all components necessary for the emergence of intracellular calcification were already present in non-calcifying ancestor cells and that no major modifications of cellular physiology and structure were required to induce intracellular calcification. An alternative and speculative hypothesis to the activation of calcium pumps could be that those pumps were already active and that the precipitation of calcium carbonate in the vesicles was prevented by inhibitors, which were subsequently lost thus leading to the emergence of intracellular calcification. But also in this alternative case, the emergence of intracellular calcification would not require major innovations in cellular structure, which is our main argument here.

3.2 Energy requirements

Since ion transport requires energy, a previous study (Monteiro et al., 2016) assumed that intracellular calcification is energetically costly, thus implying the presence of associated benefits to overcome natural selection. But because non-calcifying cells also export calcium from their cytoplasm, the energetic costs of calcification have to be quantified by comparing the costs required for ion transport into vesicles in calcifying cells versus those required by ion transport across the plasma membrane (to maintain calcium homeostasis) in both calcifying and non-calcifying cells. We therefore assessed the energetic costs of intracellular calcification by calculating the total energy required by cellular calcium transport to maintain calcium homeostasis. Our calculations indicate that the precipitation of CaCO_3 inside the vesicle removes calcium from the solution and generates a sink that depletes calcium around the cell, Eq. 2. Consequently, the gradient across the plasma membrane that drives the diffusive calcium influx through the channel is reduced with respect to non-calcifying cells, Eq. 4, thus fewer ions have to be transported out of the cytoplasm in order to maintain calcium homeostasis, Eq. 5. Since the energy required for active calcium transport out of a calcifying cell is negligibly lower than that required in non-calcifying cells (Supplementary Figure S8), we conclude that extra energy is not needed to transport calcium for intracellular calcification.




FIGURE 2

Physiological and ecological benefits of intracellular calcification. Calcifying cells (B) differ from non-calcifying cells (A) by the presence of intravesicular calcium accumulation in Golgi-derived transport vesicles (green circles represent Ca²⁺ pump/ATPase; blue cylinder indicates calcium channel). Calcium transport into vesicles allows maintenance of calcium homeostasis under elevated extracellular calcium concentrations (D) and prevents drawdown of intracellular phosphate (C) due to precipitation of Ca₃(PO₄)₂ in the cytosol. Therefore, cell division rates, black dots in (C) and (D), redrawn from Müller et al. (2015), remain unaffected by extracellular calcium concentrations in calcifying cells contrary to non-calcifying cells. The growth rate of zooplankton fed with calcifying cells (F) is generally lower than that of zooplankton fed with non-calcifying cells (E), data taken from Harvey et al. (2015) and originally produced from batch cultures in exponential growth phase by Strom et al. (2003).

However, our energy calculations also show that the reduction in energy requirements is very small because the calcification flux is slow compared to extracellular calcium diffusion so that no significant diffusive boundary layer develops at the cell surface. If all calcium ions were transported into the vesicles and no calcium were exported through the plasma membrane, $f_v = 1.0$ in Eq. 5, the energy requirements would only be 0.16 percent less than if no calcium

were transported into vesicles and all ions were exported through the plasma membrane, $f_v = 0.0$ in Eq. 5.

Besides calcium ions, also carbonate ions are needed in the vesicle to sustain calcification. The transport of carbon could thus be another process requiring energy. However, the transport of hydrogen ions that is coupled to calcium transport creates an alkaline medium inside the calcifying vesicle that shifts the




FIGURE 3

Schematic reconstruction of the steps that, based on our findings, would have led to the emergence of nannoplankton calcification. After transcription from DNA to mRNA in the nucleus, gene copies are translated into proteins in the endoplasmic reticulum. Proteins that are bound to the plasma membrane are incorporated into the ER membrane and are transported via vesicles to the Golgi apparatus. Golgi-derived vesicles are then transported to the plasma membrane where the vesicles fuse and shed their content to the extracellular space. The vesicle membrane, including all membrane proteins, becomes part of the plasma membrane. We suggest that, in a non-calcifying cell, the activation of calcium transporters occurs at the plasma membrane. Earlier activation of calcium transporters in transport vesicles would allow precipitation of sub-micron scale crystals inside the vesicles that could have led to the formation of proto-calcispheres by calcareous dinoflagellates (Janofske, 1992; Preto et al., 2013b; Dal Corso et al., 2021). The activation of calcium transporters, in the more structurally complex endomembranous systems like the Golgi apparatus, could explain the formation of delicate coccoliths and the emergence of modern-like coccolithophores. The chronological order of the appearance of calcispheres and coccolithophores shown in the diagram (from top to bottom) does not necessarily represent direct evolutionary steps regarding the activation of calcium transporters but may represent parallel origins of early Ca^{2+} transport activation that happened independently (Meier et al., 2007).

equilibrium of the carbonate system towards carbonate ions and lowers the CO_2 concentration (Holtz et al., 2013; Holtz et al., 2015). CO_2 can, therefore, passively diffuse through the vesicle membrane to replenish the carbonate ions consumed by calcification, thus sustaining calcification at moderate rates (see Figure 3 and Table 4 in Holtz et al., 2013) without additional metabolic costs. Additional carbon transport into the vesicles may only be required to sustain higher calcification rates, as observed in some coccolithophore species (e.g., *G. huxleyi*) that appeared later in the geological records. Data in support of these arguments can be found in Paasche (1962) and in Young and Ziveri (2000). Since the bicarbonate concentration in the calcifying vesicle is higher than in the cytoplasm, bicarbonate would then need to be transported actively under consumption of metabolic energy (Holtz et al., 2015). However, the bicarbonate gradient across the vesicle membrane is much smaller than the calcium gradient and CO_2 can freely diffuse across the membrane, thereby functioning as a major source of carbon to the calcifying

vesicle (Holtz et al., 2015) and, thus, preventing additional energetic requirements for bicarbonate transport.

3.3 Evolutionary success

Because of its low metabolic requirements, intracellular calcification does not require secondary evolutionary benefits to emerge. Yet, a number of hypotheses in relation to potential benefits of intracellular calcification have been proposed to explain both emergence and success of coccolithophores. These hypotheses include the potential enhancement of photosynthesis via the removal of carbonate ions (Nielsen, 1966; Paasche, 2002), protection against photoinhibition, advantages in hydrodynamic control, and protection against viruses, bacteria, and grazing (Monteiro et al., 2016). However, enhanced calcification does not enhance photosynthesis (Herfort et al., 2004; Trimborn et al., 2007)

and non-calcifying coccolithophore cells can grow even faster than calcifying cells (Rost and Riebesell, 2004). Non-calcifying strains appear also not affected by photo-inhibition at high irradiances (Nanninga and Tyrrell, 1996). Additionally, coccolithophores use non-calcifying haploid stages to escape viral infections of calcified diploid cells (Frada et al., 2008), also questioning the relevance of calcification for viral protection. All these cases cast doubts on the assumption that calcification presents relevant physiological advantages.

Previous experiments (Müller et al., 2015) showed that calcifying cells can grow faster than non-calcifying cells when exposed to elevated calcium concentrations, indicating that intracellular calcification may protect against calcium poisoning (Simkiss, 1977). Our calculations show that the Ca^{2+} -ATPase can transport ions against a gradient of up to 10^5 (see [Supplementary Material S1](#)). In non-calcifying cells, an increase of extracellular Ca^{2+} concentrations yields an increase of intracellular Ca^{2+} (Figure 2C). Since intracellular phosphate concentrations are usually around 4–5 mM (Auesukaree et al., 2004), cells operate at calcium and phosphate concentrations just below the solubility product of $\text{Ca}_3(\text{PO}_4)_2$ (Chow, 2001). Increasing intracellular calcium concentrations, therefore, lead to precipitation with phosphate due to the low solubility of $\text{Ca}_3(\text{PO}_4)_2$ (Pytkowicz and Kester, 1967; Chow, 2001). The decrease of cellular phosphate impedes cell growth in non-calcifying cells (Figure 2C). In calcifying cells, calcium transport into intracellular vesicles can still occur under elevated extracellular calcium concentrations because the calcium concentration inside the vesicle (ranging from 0.5 to 10 mM Holtz et al., 2013; Holtz et al., 2015) required for the precipitation of calcium carbonate is lower than outside the cell. The calcium gradient at the vesicle membrane therefore permits Ca^{2+} -ATPase activity and allows maintenance of calcium homeostasis even under elevated extracellular calcium concentrations (Figure 2D). Cell division of calcifying cells remains unaffected (Müller et al., 2015).

Another benefit proposed for intracellular calcification is protection against zooplankton grazing (Jaya et al., 2016; Monteiro et al., 2016). However, coccolithophores are grazed by copepods without any significant sign of saturation of ingestion rates at increasing coccolithophore densities (Harris, 1994). This indicates that calcification and thus mechanical stability does not necessarily protect against copepod grazing. Calcification can, however, have an indirect negative effect on zooplankton grazing. Culture experiments (Harvey et al., 2015) showed that ingestion of calcifying cells reduces the growth rate of the grazers (Figure 2F), whereas ingestion of non-calcifying cells leads to an increase in zooplankton growth rate (Figure 2E). A reduction of coccolithophore grazing mortality would thus provide calcifying cells with a competitive advantage against non-calcifying cells (Monteiro et al., 2016). Though not relevant for the emergence of pelagic nannoplankton calcification, the potential chemical protection against grazing may have contributed to its ecological success.

4 Conclusion

In summary, our analyses indicate that the first appearance of intracellular calcification is intrinsically linked to cellular calcium

metabolism and to the prevention of calcium accumulation in the cytoplasm. This process did not require major physiological mutations or extra energy but only a very minor change, that is the activation of calcium transport proteins during their passage from the endoplasmic reticulum to the plasma membrane. Therefore, this innovation could have occurred at any time in the geologic past because preconditions were favourable, in line with evidence suggesting unrelated appearances in different lineages. A combination of environmental factors, such as chemical conditions and ecological interactions, may have subsequently helped calcifying nannoplankton to become abundant at a global scale. Therefore, nannoplankton calcifiers rose in pelagic environments during a time, the beginning of the Mesozoic, of profound changes in the structure of Earth's ecosystems, marked by major climate changes and biological crises (i.e., the Permian–Triassic mass extinction, the Carnian Pluvial Episode and the Triassic–Jurassic mass extinction), which wiped out most of the Palaeozoic life and paved the way to the emergence of modern organisms. Within these new ecosystems, calcifying nannoplankton spread.

Data availability statement

The original contributions presented in the study are included in the article/[Supplementary Material](#), further inquiries can be directed to the corresponding author.

Author contributions

SH, JDC, and AM conceived the idea of the study. SH and ST performed the energy budget calculations. KLH compiled the genetic information and performed the gene tree analysis. All authors contributed to the interpretation of the results. SH and AM wrote the manuscript with contributions from all authors. All authors contributed to the article and approved the submitted version.

Funding

Hanse-Wissenschaftskolleg (HWK) provided visiting fellowships to JDC and KLH.

Acknowledgments

We gratefully acknowledge the Hanse-Wissenschaftskolleg (HWK), Delmenhorst, Germany, for providing visiting fellowship to JDC and KLH and thus for creating the opportunity to bring together the multidisciplinary expertise necessary for this study.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships

that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be

evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/feart.2023.747059/full#supplementary-material>

References

- Aueskaree, C., Homma, T., Tochio, H., Shirakawa, M., Kaneko, Y., and Harashima, S. (2004). Intracellular phosphate serves as a signal for the regulation of the PHO pathway in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 279, 17289–17294. doi:10.1074/jbc.M312202200
- Benner, I., Diner, R. E., Lefebvre, S. C., Li, D., Komada, T., Carpenter, E. J., et al. (2013). *Emiliana huxleyi* increases calcification but not expression of calcification-related genes in long-term exposure to elevated temperature and pCO₂. *Philosophical Trans. R. Soc. B* 368, 20130049. doi:10.1098/rstb.2013.0049
- Berner, R. A., and Kothavala, Z. (2001). Geocarb III: A revised model of atmospheric CO₂ over phanerozoic time. *Am. J. Sci.* 301, 182–204. doi:10.2475/ajs.301.2.182
- Bown, P. R. (1998). *Calcareous nannofossil biostratigraphy*. Dordrecht: Springer.
- Bown, P. R., Lees, J. A., and Young, J. R. (2004). "Calcareous nannoplankton evolution and diversity through time," in *Coccolithophores. From molecular processes to global impact* (Springer Berlin), 481–508.
- Boynton, A. L., Whitfield, F. F., and MacManus, J. P. (1980). Calmodulin stimulates DNA synthesis by Rat Liver cells. *Biochem. Biophysical Res. Commun.* 95, 745–749. doi:10.1016/0006-291x(80)90849-9
- Bralower, T. J., Bown, P. R., and Siesser, W. G. (1991). Significance of upper triassic nannofossils from the southern hemisphere (ODP leg 122, wombat plateau, NW Australia). *Mar. Micropaleontol.* 17, 119–154. doi:10.1016/0377-8398(91)90025-2
- Brownlee, C., Davies, M., Nimer, N., Dong, L., and Merrett, M. (1995). Calcification, photosynthesis and intracellular regulation in *Emiliana huxleyi*. *Bull. l'institute Océanogr. Monaco* 14, 19–36.
- Brownlee, C., Wheeler, G. L., and Taylor, A. R. (2015). Coccolithophore biomineralization: New questions, new answers. *Seminars Cell and Dev. Biol.* 46, 11–16. doi:10.1016/j.semcdb.2015.10.027
- Carafoli, E. (1987). Intracellular calcium homeostasis. *Ann. Rev. Biochem.* 56, 395–433. doi:10.1146/annurev.bi.56.070187.002143
- Chow, L. (2001). Solubility of calcium phosphates. *Octacalcium Phosphate* 1, 94–111. doi:10.1159/000061650
- Dal Corso, J., Preto, N., Agnini, C., Hohn, S., Merico, A., Willems, H., et al. (2021). Rise of calcispheres during the carnian pluvial Episode (late triassic). *Glob. Planet. Change* 200, 103453. doi:10.1016/j.gloplacha.2021.103453
- de Vargas, C., Aubry, M. P., Probert, I., and Young, J. (2007). Origin and evolution of coccolithophores. From coastal hunters to oceanic farmers. *Evol. Prim. Prod. Sea* 2007, 251–285. doi:10.1016/B978-012370518-1/50013-8
- Demangel, I., Kovács, Z., Richoz, S., Gardin, S., Krystyn, L., Baldermann, A., et al. (2020). Development of early calcareous nannoplankton in the late triassic (northern calcareous alps, Austria). *Glob. Planet. Change* 193, 103254. doi:10.1016/j.gloplacha.2020.103254
- Frada, M., Probert, I., Allen, M. J., Wilson, W. H., and de Vargas, C. (2008). The Cheshire Cat escape strategy of the coccolithophore *Emiliana huxleyi* in response to viral infection. *Proc. Natl. Acad. Sci.* 105, 15944–15949. doi:10.1073/pnas.0807707105
- Gardin, S., Krystyn, L., Richoz, S., Bartolini, A., and Galbrun, B. (2012). Where and when the earliest coccolithophores? *Lethaia* 45, 507–523. doi:10.1111/j.1502-3931.2012.00311.x
- Goddéris, Y., Donnadieu, Y., Vargas, C. D., Pierrehumbert, R. T., Dromart, G., and van de Schootbrugge, B. (2008). Causal or casual link between the rise of nannoplankton calcification and a tectonically-driven massive decrease in Late Triassic atmospheric CO₂? *Earth Planet. Sci. Lett.* 267, 247–255. doi:10.1016/j.epsl.2007.11.051
- Gussone, N., Langer, G., Thoms, S., Nehrke, G., Eisenhauer, A., Riebesell, U., et al. (2006). Cellular calcium pathways and isotope fractionation in *Emiliana huxleyi*. *Geology* 34, 625–628. doi:10.1130/G22733.1
- Harris, R. P. (1994). Zooplankton grazing on the coccolithophore *Emiliana huxleyi* and its role in inorganic carbon flux. *Mar. Biol.* 119, 431–439. doi:10.1007/BF00347540
- Harvey, E. L., Bidle, K. D., and Johnson, M. D. (2015). Consequences of strain variability and calcification in *Emiliana huxleyi* on microzooplankton grazing. *J. Plankton Res.* 37, fbv081–1148. doi:10.1093/plankt/fbv081
- Hepler, P. K. (1994). The role of calcium in cell division. *Cell Calcium* 16, 322–330. doi:10.1016/0143-4160(94)90096-5
- Herfort, L., Loste, E., Meldrum, F., and Thake, B. (2004). Structural and physiological effects of calcium and magnesium in *Emiliana huxleyi* (Lohmann) Hay and Mohler. *J. Struct. Biol.* 148, 307–314. doi:10.1016/j.jsb.2004.07.005
- Holtz, L. M., Thoms, S., Langer, G., and Wolf-Gladrow, D. A. (2013). Substrate supply for calcite precipitation in *Emiliana huxleyi*: Assessment of different model approaches. *J. Phycol.* 49, 417–426. doi:10.1111/jpy.12052
- Holtz, L. M., Wolf-Gladrow, D., and Thoms, S. (2015). Numerical cell model investigating cellular carbon fluxes in *Emiliana huxleyi*. *J. Theor. Biol.* 364, 305–315. doi:10.1016/j.jtbi.2014.08.040
- Inesi, G., and Tadini-Buoninsegni, F. (2014). Ca²⁺/H⁺ exchange, luminal Ca²⁺ release and Ca²⁺/ATP coupling ratios in the sarcoplasmic reticulum ATPase. *J. Cell Commun. Signal.* 8, 5–11. doi:10.1007/s12079-013-0213-7
- Inouye, I., and Pienaar, R. N. (1983). Observations on the life cycle and microanatomy of *Thoracosphaera heimii* (Dinophyceae) with special reference to its systematic position. *S. Afr. Tydskr. Plantk.* 2, 63–75. doi:10.1016/S0022-4618(16)30147-4
- Janofske, D. (1992). Kalkiges Nannoplankton, insbesondere Kalkige Dinoflagellaten-Zysten der alpinen Ober-Trias: Taxonomie, Biostratigraphie und Bedeutung für die Phylogenie der Peridinales. *Berl. Geowiss. Abh. Reihe E* 4, 53.
- Jantschke, A., Pinkas, I., Schertel, A., Addadi, L., and Weiner, S. (2020). Biomineralization pathways in calcifying dinoflagellates: Uptake, storage in mgcap-rich bodies and formation of the shell. *Acta Biomater.* 102, 427–439. doi:10.1016/j.actbio.2019.11.042
- Jaya, B. N., Hoffmann, R., Kirchlechner, C., Dehm, G., Scheu, C., and Langer, G. (2016). Coccospheeres confer mechanical protection: New evidence for an old hypothesis. *Acta Biomater.* 42, 258–264. doi:10.1016/j.actbio.2016.07.036
- Kaźmierczak, J., Ittekkot, V., and Degens, E. T. (1985). Biocalcification through time: Environmental challenge and cellular response. *Paläontologische Z.* 59, 15–33. doi:10.1007/BF02985996
- Kaźmierczak, J., Kempe, (2004). "Calcium build-up in the precambrian sea," in *Origins: Genesis, evolution and diversity of life* (Springer), 329–345.
- Kaźmierczak, J., and Kremer, B. (2005). Early post-mortem calcified Devonian acritarchs as a source of calcispheric structures. *Facies* 51, 554–565. doi:10.1007/s10347-005-0071-8
- Kearse, M., Moir, R., Wilson, A., Stones-havas, S., Cheung, M., Sturrock, S., et al. (2012). Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28, 1647–1649. doi:10.1093/bioinformatics/bts199
- Knoll, A. H. (2003). Biomineralization and evolutionary history. *Rev. Mineralogy Geochem.* 54, 329–356. doi:10.2113/0540329
- Krishtal, O. A., Pidoplichko, V. I., and Shakhvalov, Y. A. (1981). Conductance of the calcium channel in the membrane of snail neurones. *J. physiology* 310, 423–434. doi:10.1113/jphysiol.1981.sp013558
- Langer, G., De Nooijer, L. J., and Oetjen, K. (2010). On the role of the cytoskeleton in coccolith morphogenesis: The effect of cytoskeleton inhibitors. *J. Phycol.* 46, 1252–1256. doi:10.1111/j.1529-8817.2010.00916.x
- Langer, G., Gussone, N., Nehrke, G., Riebesell, U., Eisenhauer, A., Kuhnert, H., et al. (2006). Coccolith strontium to calcium ratios in *Emiliana huxleyi*: The dependence on seawater strontium and calcium concentrations. *Limnol. Oceanogr.* 51, 310–320. doi:10.4319/lo.2006.51.1.0310

- Lewis, E., and Wallace, D. (1998). *CO2SYS program. Tech. Rep., carbon dioxide information analysis center*. Oak Ridge, Tennessee: Oak Ridge National Laboratory Environmental Sciences Division.
- Lowenstam, H. A., and Margulis, L. (1980). Evolutionary prerequisites for early Phanerozoic calcareous skeletons. *BioSystems* 12, 27–41. doi:10.1016/0303-2647(80)90036-2
- Lux, H. D., and Nagy, K. (1981). Single Channel Ca^{2+} currents in helix pomatia neurons. *Pflügers Archiv Eur. J. Physiology* 391, 252–254. doi:10.1007/bf00596179
- Marcum, J. M., Dedman, J. R., Brinkley, B. R., and Means, A. R. (1978). Control of microtubule assembly-disassembly by calcium-dependent regulator protein. *Proc. Natl. Acad. Sci.* 75, 3771–3775. doi:10.1073/pnas.75.8.3771
- Meier, K. J., Young, J. R., Kirsch, M., and Feist-Burkhardt, S. (2007). Evolution of different life-cycle strategies in oceanic calcareous dinoflagellates. *Eur. J. Phycol.* 42, 81–89. doi:10.1080/09670260600937833
- Meldolesi, J., and Pozzan, T. (1998). The endoplasmic reticulum Ca^{2+} store: A view from the lumen. *Trends Biochem. Sci.* 23, 10–14. doi:10.1016/S0968-0004(97)01143-2
- Montañez, I. P., McElwain, J. C., Poulsen, C. J., White, J. D., DiMichele, W. A., Wilson, J. P., et al. (2016). Climate, p_{CO_2} and terrestrial carbon cycle linkages during late palaeozoic glacial–interglacial cycles. *Nat. Geosci.* 9, 824–828. doi:10.1038/ngeo2822
- Monteiro, F. M., Bach, L. T., Brownlee, C., Bown, P., Rickaby, R. E. M., Poulton, A. J., et al. (2016). Why marine phytoplankton calcify. *Sci. Adv.* 2, e1501822. doi:10.1126/sciadv.1501822
- Müller, M. N., Barcelos E Ramos, J., Schulz, K. G., Riebesell, U., Kaźmierczak, J., Gallo, F., et al. (2015). Phytoplankton calcification as an effective mechanism to alleviate cellular calcium poisoning. *Biogeosciences* 12, 6493–6501. doi:10.5194/bg-12-6493-2015
- Nanninga, H. J., and Tyrrell, T. (1996). Importance of light for the formation of algal blooms by *Emiliania huxleyi*. *Mar. Ecol. Prog. Ser.* 136, 195–203. doi:10.3354/meps136195
- Nielsen, E. S. (1966). The uptake of free CO_2 and HCO_3^- during photosynthesis of plankton algae with special reference to the coccolithophorid *coccolithus huxleyi*. *Physiol. Plant.* 19, 232–240. doi:10.1111/j.1399-3054.1966.tb09095.x
- Onoue, T., and Yoshida, A. (2010). Depositional response to the Late Triassic ascent of calcareous plankton in pelagic mid-oceanic plate deposits of Japan. *J. Asian Earth Sci.* 37, 312–321. doi:10.1016/j.jseas.2009.08.013
- Orrenius, S., McConkey, D. J., Bellomo, G., and Nicotera, P. (1989). Role of Ca^{2+} in toxic cell killing. *Trends Pharmacol. Sci.* 10, 281–285. doi:10.1016/0165-6147(89)90029-1
- Paasche, E. (2002). A review of the coccolithophorid *Emiliania huxleyi* (Prymnesiophyceae), with particular reference to growth, coccolith formation, and calcification-photosynthesis interactions. *Phycologia* 40, 503–529. doi:10.2216/i0031-8884-40-6-503.1
- Paasche, E. (1962). Coccolith Formation. *Nature* 193, 1094–1095. doi:10.1038/1931094b0
- Pfeffer, S. R., and Rothmann, J. E. (1987). Biosynthetic protein transport and sorting by the endoplasmic reticulum and golgi. *Annu. Rev. Biochem.* 56, 829–852. doi:10.1146/annurev.bi.56.070187.004145
- Poenie, M., Alderton, J., Tsien, R. Y., and Steinhardt, R. A. (1985). Changes of free calcium levels with stages of the cell division cycle. *Nature* 315, 147–149. doi:10.1038/315147a0
- Preto, N., Agnini, C., Rigo, M., Sprovieri, M., and Westphal, H. (2013a). The calcareous nannofossil *Prinsiosphaera* achieved rock-forming abundances in the latest Triassic of Western Tethys: Consequences for the $\delta^{13}C$ of bulk carbonate. *Biogeosciences* 10, 6053–6068. doi:10.5194/bg-10-6053-2013
- Preto, N., Willems, H., Guaiumi, C., and Westphal, H. (2013b). Onset of significant pelagic carbonate accumulation after the carnian pluvial event (CPE) in the Western tethys. *Facies* 59, 891–914. doi:10.1007/s10347-012-0338-9
- Pytkowicz, R. M., and Kester, D. R. (1967). Relative calcium phosphate saturation in two regions of the north pacific ocean. *Limnol. Oceanogr.* 12, 714–718. doi:10.4319/lo.1967.12.4.0714
- Raup, D., and Sepkoski, J., Jr (1982). Mass extinctions in the marine fossil record. *Science* 215, 1501–1503. doi:10.1126/science.215.4539.1501
- Read, B. A., Kegel, J., Klute, M. J., Kuo, A., Lefebvre, S. C., Maumus, F., et al. (2013). Pan genome of the phytoplankton *Emiliania* underpins its global distribution. *Nature* 499, 209–213. doi:10.1038/nature12221
- Ridgwell, A. (2005). A Mid Mesozoic Revolution in the regulation of ocean chemistry. *Mar. Geol.* 217, 339–357. doi:10.1016/j.margeo.2004.10.036
- Ridgwell, A., and Zeebe, R. E. (2005). The role of the global carbonate cycle in the regulation and evolution of the Earth system. *Earth Planet. Sci. Lett.* 234, 299–315. doi:10.1016/j.epsl.2005.03.006
- Rost, B., and Riebesell, U. (2004). “Coccolithophores and the biological pump: Responses to environmental changes,” in *Coccolithophores: From molecular processes to global impact*. Editors H. R. Thierstein, and J. R. Young (Springer Berlin), 99–125. doi:10.1007/978-3-662-06278-4_5
- Royer, D. L., Berner, R. A., Montañez, I. P., Tabor, N. J., and Beerling, D. J. (2004). CO_2 as a primary driver of Phanerozoic climate. *Geol. Soc. Am. Today* 14, 4–10. doi:10.1130/1052-5173(2004)014<4:caapdo>2.0.co;2
- Sanders, D., Pelloux, J., Brownlee, C., and Harper, J. F. (2002). Calcium at the crossroads of signaling. *Plant Cell* 14 (1), S401–S417. doi:10.1105/tpc.002899
- Sepkoski, J. J., and Ausich, W. I. (1990). “Evolutionary faunas,” in *Palaeobiology: A synthesis*. Editors D. E. G. Briggs, and P. R. Crowther (Oxford: Blackwell Scientific Publications), 37–41.
- Simkiss, K. (1977). Biomineralization and detoxification. *Calcif. Tissue Res.* 24, 199–200. doi:10.1007/BF0222316
- Stanley, S. M., and Hardie, L. A. (1998). Secular oscillations in the carbonate mineralogy of reef-building and sediment-producing organisms driven by tectonically forced shifts in seawater chemistry. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 144, 3–19. doi:10.1016/S0031-0182(98)00109-6
- Steinhardt, R. A., and Alderton, J. (1988). Intracellular free calcium rise triggers nuclear envelope breakdown in the sea urchin embryo. *Nature* 332, 364–366. doi:10.1038/332364a0
- Strom, S., Wolfe, G., Holmes, J., Stecher, H., Shimeneck, C., and Sarah, L. (2003). Chemical defense in the microplankton i: Feeding and growth rates of heterotrophic protists on the dms-producing phytoplankton *emiliania huxleyi*. *Limnol. Oceanogr.* 48, 217–229. doi:10.4319/lo.2003.48.1.0217
- Tidow, H., Poulsen, L. R., Andreeva, A., Knudsen, M., Hein, K. L., Wiuf, C., et al. (2012). A bimodular mechanism of calcium control in eukaryotes. *Nature* 491, 468–472. doi:10.1038/nature11539
- Toyoshima, C., and Inesi, G. (2004). Structural basis of ion pumping by Ca^{2+} -ATPase of the sarcoplasmic reticulum. *Annu. Rev. Biochem.* 73, 269–292. doi:10.1146/annurev.biochem.73.011303.073700
- Trimborn, S., Langer, G., and Rost, B. (2007). Effect of varying calcium concentrations and light intensities on calcification and photosynthesis in *Emiliania huxleyi*. *Limnol. Oceanogr.* 52, 2285–2293. doi:10.4319/lo.2007.52.5.2285
- Tsien, R. W. (1983). Calcium channels in excitable cell membranes. *Annu. Rev. Physiology* 45, 341–358. doi:10.1146/annurev.ph.45.030183.002013
- Verret, F., Wheeler, G., Taylor, A. R., Farnham, G., and Brownlee, C. (2010). Calcium channels in photosynthetic eukaryotes: Implications for evolution of calcium based signalling. *New Phytol.* 187, 23–43. doi:10.1111/j.1469-8137.2010.03271.x
- von Dassow, P., Ogata, H., Probert, I., Wincker, P., Da Silva, C., Audic, S., et al. (2009). Transcriptome analysis of functional differentiation between haploid and diploid cells of *Emiliania huxleyi*, a globally significant photosynthetic calcifying cell. *Genome Biol.* 10, R114. doi:10.1186/gb-2009-10-10-r114
- Winther, A.-M. L., Bublitz, M., Karlens, J. L., Møller, J. V., Hansen, J. B., Nissen, P., et al. (2013). The sarcolipin-bound calcium pump stabilizes calcium sites exposed to the cytoplasm. *Nature* 495, 265–269. doi:10.1038/nature11900
- Witkowski, C. R., Weijers, J. W., Blais, B., Schouten, S., and Sinninghe Damsté, J. S. (2018). Molecular fossils from phytoplankton reveal secular P_{CO_2} trend over the Phanerozoic. *Sci. Adv.* 4, eaat4556. doi:10.1126/sciadv.aat4556
- Young, J. R. (2003). Biomineralization within vesicles: The calcite of coccoliths. *Rev. Mineralogy Geochem.* 54, 189–215. doi:10.2113/0540189
- Young, J. R., Geisen, M. P. I., and Probert, I. (2005). A review of selected aspects of coccolithophore biology with implications for paleobiodiversity estimation. *Micropaleontology* 51, 267–288. doi:10.2113/gsmicropal.51.4.267
- Young, J. R., and Ziveri, P. (2000). Calculation of coccolith volume and its use in calibration of carbonate flux estimates. *Deep-Sea Res. Part II Top. Stud. Oceanogr.* 47, 1679–1700. doi:10.1016/S0967-0645(00)00003-5
- Zeebe, R. E., and Westbroek, P. (2003). A simple model for the $CaCO_3$ saturation state of the ocean: The Strangelove, the Neritan, and the Cretan Ocean. *Geochem. Geophys. Geosystems* 4. doi:10.1029/2003GC000538