

# **Maps of the Geographical Distribution of Macrozooplankton in the Atlantic Sector of the Southern Ocean**

by Uwe Piatkowski

**Berichte zur Polarforschung Nr. 22 /Februar 1985**  
**Reports on Polar Research no. 22 /February 1985**

---



Contents

	page
Summary - Zusammenfassung .....	4
1. Introduction .....	5
2. Material and Methods .....	5
3. Results .....	8
4. References .....	12
5. Maps of Macrozooplankton Distribution .....	15

### Summary

Macrozooplankton samples were taken during three cruises of the FRV "Walther Herwig" into the Atlantic sector of the Southern Ocean during the austral summers of 1975/76, 1977/78 and 1980/81.

The present paper presents maps on the geographical distribution and abundance of the five most frequent macrozooplankton groups in the Southern Ocean, adult krill (Euphausia superba), other euphausiids, amphipods, chaetognaths and salps.

### Zusammenfassung

Während der drei Forschungsreisen des FFS "Walther Herwig" in den atlantischen Sektor des Südpolarmeeres, die in den Südsommern 1975/76, 1977/78 und 1980/81 stattfanden, wurde ein umfangreiches Makrozooplankton-Material gesammelt.

Der vorliegende Bericht untersucht die geographische Verbreitung und relative Häufigkeit von adultem Krill (Euphausia superba), anderen Euphausiiden, Amphipoden, Chaetognathen und Salpen, die die fünf häufigsten Makrozooplankton-Gruppen im Südpolarmeer darstellen.

# MAPS OF THE GEOGRAPHICAL DISTRIBUTION OF MACROZOOPLANKTON IN THE ATLANTIC SECTOR OF THE SOUTHERN OCEAN

by Uwe Piatkowski

## 1. Introduction

The main purpose of the German Antarctic Expeditions with FRV "Walther Herwig" into the Atlantic sector of the Southern Ocean was research on Antarctic krill (Euphausia superba) and commercial fish species.

Sahrhage et al. (1978), Hempel et al. (1979) and Hempel (1982) have reported comprehensively on the three expeditions, which took place during the austral summers of 1975/76, 1977/78 and 1980/81.

Net sampling programmes were carried out to investigate distribution and abundance of krill and to analyse population structures. Sampling gear consisted of a pelagic midwater trawl, a Bonqo net and a rectangular midwater trawl (RMT 1+8) (Baker et al. 1973). The latter was used in conjunction with echo-surveys.

The krill samples obtained were studied intensively by several authors (i.g. Nast 1978, 1982a, b, Nast et al. 1982, Pommeranz 1978a, b, Siegel 1982, 1983, Wörner 1979).


Macrozooplankton groups other than krill that were examined, included other euphausiids (Weidmann-Haass and Haass 1980), amphipods (Andres 1978, 1982, 1983, Weidmann-Haass 1983), chaetognaths (James 1979), postlarval notothenioids (Kellermann and Kock 1984) and myctophids (Rowedder 1979).

This report is part of a master thesis (Piatkowski 1982). It describes the geographical distribution and abundance of the five prevailing macrozooplankton taxa caught with the RMT 8 on all three expeditions. They are krill (Euphausia superba), other euphausiids, amphipods, chaetognaths and salps. The distribution charts of Euphausia superba, shown in this study, are modified figures previously published by Pommeranz (1978), Wörner (1979), Nast (1982) and Siegel (1982).

## 2. Material and Methods

Samples were taken in the Atlantic sector of the Southern Ocean between 30°W and 64°W. The area surveyed was subdivided into three regions (Fig. 1):

Fig. 1: Subdivisions of the area investigated.



South Shetland Islands  
Central Scotia Sea  
South Georgia.

All macrozooplankton samples analysed were obtained with the RMT 8 (mesh size 4.5 mm, mouth opening approximately 8 m<sup>2</sup>) of the Institute of Oceanographic Science's RMT 1+8 (Baker et al. 1973). The standard procedure was an oblique haul in the surface waters (maximum fishing depth 200 m).

Volume of filtered water was calculated by Pommeranz (unpublished). Station maps and lists, detailed descriptions of hauls, accurate haul depths and comments on stations were published by

Wörner and Kühn (1978) for cruise 1975/76  
Wörner (1978) for cruise 1977/78  
Piatkowski and Klages (1983) for cruise 1980/81.

Sorting procedures and results of the micronekton and zooplankton investigations were given by

James and Wörner (1978) for cruise 1975/76  
James (1980) for cruise 1977/78  
Piatkowski et al. (1984) for cruise 1980/81.

James and Wörner (1978), assisted by several students sorted the RMT 8 samples into the following taxa:

Fish  
Fish larvae  
Fish eggs  
Scyphomedusae  
Euphausia superba (postlarvae and adults)  
Other Euphausiids (postlarvae and adults)  
Amphipods  
Decapods (postlarvae and adults)  
Mysids  
Chaetognaths  
Salps  
Siphonophores  
Polychaetes  
Cephalopods  
Other Molluscs  
Copepods + Ostracods (presence/absence).

For this study only Euphausia superba, other euphausiids, amphipods, chaetognaths, and salps were considered for analyses of geographical distribution and abundance, as they were clearly the most abundant macrozooplankton groups.

### 3. Results

A total of 357 samples was analysed in this study. In Table 1 they are compiled according to geographical locations and time of year.

#### Composition of macrozooplankton groups

Euphausia superba was the prevailing macrozooplankton species in all three areas surveyed. Other euphausiids found, were mainly Thysanoessa macrura, E. frigida and E. triacantha, of which T. macrura was the most abundant euphausiid next to E. superba. The major representative amphipod was the hyperiid Themisto gaudichaudii. Of the chaetognaths, the species Eukrohnia hamata and Sagitta gazellae prevailed. Salpa thompsoni was the absolute dominant species of the salps.

#### Geographical distribution and abundance

Table 2 presents the density classes, in which the major macrozooplankton taxa occurred. Only E. superba and salps were found in densities higher than 10000 ind/m<sup>2</sup>.

Maximum concentrations of different macrozooplankton groups in each of the three regions are shown in Table 3:

Table 3 Maximum concentrations of different macrozooplankton groups in the three regions investigated (individuals/m<sup>2</sup>).

Group	South Shetland Islands	Central Scotia Sea	South Georgia
<u>Euphausia superba</u>	7766	1727	333746
Other euphausiids	1074	143	667
Amphipods	168	380	505
Chaetognaths	97	511	1053
Salps	36033	41520	1896

Figures 2 - 81 show the geographical distribution and abundance of the predominant taxa in the three regions investigated.

Each figure represents a different month and cruise leg, thus monthly and annual differences in the distribution and abundance of the macrozooplankton groups in each of the three regions become apparent.



Table 1 Number of analysed RMT 8 samples compiled according to geographical locations and time of year.

Location	November	December	January	February	March	April	Total
<b>South Shetland Islands</b>							
Cruise 1975/76	-	-	4	17	-	-	21
Cruise 1977/78	27	12	16	8	42	-	105
Cruise 1980/81	-	-	3	6	23	-	32
<b>Central Scotia Sea</b>							
Cruise 1975/76	4	20	1	11	7	-	43
Cruise 1977/78	7	7	9	5	6	-	34
Cruise 1980/81	-	-	-	24	-	-	24
<b>South Georgia</b>							
Cruise 1975/76	-	34	-	-	1	7	42
Cruise 1977/78	16	8	9	-	17	6	56
Cruise 1980/81	-	-	-	-	-	-	-

Table 2 Number and percentage of catches separated in major macrozooplankton taxa and grouped in density classes.

Density class Individuals/m <sup>2</sup>	Euphausia number	superba %	Other euphausiids number	%	Amphipods number	%	Chaetognaths number	%	Salps number	%
≤ 0.1	59	22.5	19	5.8	57	17.2	51	17.3	33	13.9
0.1 - 1	87	33.2	84	25.5	120	36.3	120	40.7	64	26.9
1.1 - 10	47	17.9	141	42.7	113	34.1	105	35.6	66	27.7
11 - 100	44	16.8	74	22.4	34	10.3	16	5.4	41	17.2
101 - 1000	17	6.5	11	3.3	7	2.1	2	0.7	19	8.0
1001 - 10000	6	2.3	1	0.3	0	0	1	0.3	11	4.6
≥ 10000	2	0.8	0	0	0	0	0	0	4	1.7
Positive catches	262	74.3	330	92.0	331	92.8	295	82.7	238	67.2
Zero catches	95	25.7	27	8.0	26	7.2	62	17.3	119	32.8
Total	357	100.0	357	100.0	357	100.0	357	100.0	357	100.0

Particularly rich samples of Euphausia superba ( $\geq 1000$  ind/m<sup>2</sup>) were found in the waters off Elephant Island, in the north-western part of the Weddell Sea and on the northern shelf regions of South Georgia. In November/December krill concentrations were significantly higher than in the following months. E. superba distribution was heterogeneous. Apart from maximum concentrations of all groups investigated E. superba also showed the highest number of samples with densities less than 0.1 ind/m<sup>2</sup>.

Other euphausiids, amphipods and chaetognaths occurred in distinctly lower numbers than E. superba and salps. They had a more even distribution than E. superba in all areas investigated. Annual or monthly differences in abundance could not be detected.

Highest concentrations of salps were obtained during January and February in all regions. The numbers caught were significantly high, if E. superba occurred in low densities (Figs. 36 and 56, 57 and 77).

Comparing the abundances of different times of the day only salps indicated patterns of a diurnal migration, because their numbers increased in the surface waters during the hauls at dusk.

The data did not reveal a significant diurnal migration of the other groups, these appeared to remain within the surface layers ( $\leq 200$  m water depth) both during the day and night (Table 4).

Table 4 Mean abundances of investigated macrozooplankton groups (individuals/m<sup>2</sup>) at various times of the day. DA = daytime, DU = dusk, NI = nighttime, DW = dawn.

Group	DA	DU	NI	DW
<u>Euphausia superba</u>	1426	21	519	74
Other euphausiids	20	10	15	14
Amphipods	11	5	6	2
Chaetognaths	10	$\leq 1$	2	1
Salps	492	2188	127	11

The abundance of all five groups, especially that of the salps was significantly higher in the season of 1975/76 than in the seasons of 1977/78 and 1980/81, which is applicable to all three regions investigated.

#### Acknowledgements

I thank all scientists as well as the crew of the FRV "Walther Herwig", who were engaged in the plankton sampling programmes during the three cruises. I am grateful to the many colleagues, who carefully sorted the numerous plankton samples. Dr. T. Pommeranz kindly supplied his unpublished data of the filtered water volumes of the RMT 8. Thanks are due to Dr. G. Dieckmann for his help in correcting the English and to Mrs. S. Marschall for typing the draft.

#### 4. References

- Andres HG (1978) Gammaridea (Amphipods, Crustacea) der Antarktis-Expedition 1975/76. Auswertung der Dauerstation südlich von Elephant Island. Meeresforsch 27: 88-102
- Andres HG (1982) Die Gammaridea (Crustacea: Amphipoda) der Deutschen Antarktis-Expeditionen 1975/76 und 1977/78. 2. Eusiridae. Mitt hamb zool Mus Inst 79: 159-185
- Andres HG (1983) Die Gammaridea (Crustacea: Amphipoda) der Deutschen Antarktisexpeditionen 1975/76 und 1977/78. 3. Lysianassidae. Mitt hamb zool Mus Inst 80: 183-220
- Baker A de C, Clarke MR, Harris MJ (1973) The N.I.O. Combination Net (RMT 1+8) and further developments of Rectangular Midwater Trawls. J mar biol Ass UK 53: 167-184
- Hempel G (1982) Antarktis-Expedition 1981 der Bundesrepublik Deutschland mit FFS "Walther Herwig". Einleitung, Aufgabe der Expedition, Organisation sowie nationale und internationale Zusammenarbeit. Arch FischWiss 33 (Beih 1): 5-15
- Hempel G, Sahrhage D, Schreiber W, Steinberg R (1979) Antarktis-Expedition 1977/78 der Bundesrepublik Deutschland. Arch FischWiss 30 (Beih 1), pp 1-119
- James R (1979) Antarctic chaetognaths from a time station near Elephant Island. Meeresforsch 27: 282-287
- James R (1980) Sortiererergebnisse des Zooplanktons der deutschen Antarktis-Expedition 1977/78. Schriftl Mitt, pp 1-50
- James R, Wörner FG (1978) Results of the sorting of the mikronekton and zooplankton material sampled by the German Antarctic Expedition 1975/76. Ber Inst Meereskd Kiel 59, pp 1-53

- Kellermann A, Kock K-H (1984) Postlarval and Juvenile Notothenioids (Pisces, Perciformes) in the Southern Scotia Sea and Northern Weddell Sea during FIBEX 1981. *Meeresforsch* 30: 82-93
- Nast F (1978) The vertical distribution of larval and adult krill (*Euphausia superba* Dana) on a time station south of Elephant Island, South Shetlands. *Meeresforsch* 27: 103-118
- Nast F (1982a) Krillfänge während FIBEX 1981. *Arch FischWiss* 33 (Beih 1): 61-84
- Nast F (1982b) The assessment of krill (*Euphausia superba* Dana) biomass from a net sampling programme. *Meeresforsch* 29: 154-165
- Nast F, Pommeranz T, Stein M (1982) A comparison of krill abundance between the 1975/76 and 1980/81 seasons around the Weddell-Scotia confluence. *ICES Biol Oceanogr Cttee CM L:5*, pp 1-11
- Piatkowski U (1982) Beiträge zur Verbreitung des Makrozooplanktons in der Scotia Sea und ihrer angrenzenden Gebiete. Masters Thesis Univ Kiel, pp 1-149
- Piatkowski U, Klages N (1983) German Antarctic Expedition 1980/81 with FRV "Walther Herwig" and RV "Meteor". First International BIOMASS Experiment (FIBEX). Data of micronekton and zooplankton hauls. *Ber Polarforsch* 15, pp 1-57
- Piatkowski U, Hempel I, Rakusa-Suszczewski S (1984) FIBEX Cruise Zooplankton Data. *Ber Polarforsch* 16, pp 1-53
- Pommeranz T (1978a) Mikronekton und Zooplankton. In: Sahrhage D, Schreiber W, Steinberg R, Hempel G (eds) *Antarktische Expedition 1975/76 der Bundesrepublik Deutschland*. *Arch FischWiss* 29 (Beih 1): 31-41
- Pommeranz T (1978b) Attempts for Quantitative Studies in the Distribution of Adult Krill *Euphausia superba* Dana in the Scotia Sea. *ICES Biol Oceanogr Cttee CM L:5*, pp 1-17
- Rowedder U (1979) Some aspects of the biology of *Electrona antarctica* (Günther, 1878), (Family Myctophidae). *Meeresforsch* 27: 244-251
- Sahrhage D, Schreiber W, Steinberg R, Hempel G (1978) *Antarktische Expedition 1975/76 der Bundesrepublik Deutschland*. *Arch FischWiss* 29 (Beih 1), pp 1-96

- Siegel V (1982) Untersuchungen an Nachlaichkonzentrationen des antarktischen Krills Euphausia superba. Arch FischWiss 33 (Beih 1): 113-125
- Siegel V (1983) Population structure of Euphausia superba in the eastern part of the Bransfield Strait. In: Schnack SB (ed) On the Biology of Krill (Euphausia superba). Proceedings of the Seminar and Report of the Krill Biology Group, Bremerhaven, 12-16 May 1983. Ber Polarforsch Sonderh 4: 227-238
- Weigmann-Haass R (1983) Zur Taxonomie und Verbreitung der Gattung Cylopus Dana 1853 (Amphipoda:Hyperidei) im antarktischen Teil des Atlantik. Meteor Forsch Ergebnisse (Reihe D) 36: 1-11
- Weigmann-Haass R, Haass G (1980) Geographische Verbreitung und vertikale Verteilung der Euphausiacea (Crustacea) während der Antarktis-Expedition 1975/76. Meeresforsch 28: 19-31
- Wörner FG (1978) Liste der Mikronekton- und Zooplanktonfänge der 2. Deutschen Antarktis-Expedition 1977/78. Ber Inst Meereskd Kiel 60, pp 1-82
- Wörner FG (1979) Zooplankton- und Mikronektonfänge. In: Hempel G, Sahrhage D, Schreiber W, Steinberg R (eds) Antarktis-Expedition 1977/78 der Bundesrepublik Deutschland. Arch FischWiss 30 (Beih 1): 40-61
- Wörner FG, Kühn A (1978) Liste der Mikronekton- und Zooplanktonfänge der deutschen Antarktis-Expedition 1975/76. Ber Inst Meereskd Kiel 48, pp 1-55

5. Maps of Macrozooplankton Distribution

South Shetland Islands




Fig. 2: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in November/December 1977 (modified from Wörner 1979).




Fig. 3: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in January/February 1976 (modified from Pommeranz 1978).




Fig. 4: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in January/February 1978 (modified from Wörner 1979).




Fig. 5: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in January/February 1981 (modified from Nast 1982b).






Fig. 6: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in March 1978 (modified from Wörner 1979).




Fig. 7: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in March 1981 (modified from Siegel 1982).




Fig. 8: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in November/December 1977.




Fig. 9: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in January/February 1976.




Fig. 10: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in January/February 1978.




Fig. 11: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in January/February 1981.




Fig. 12: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in March 1978.




Fig. 13: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in March 1981.




Fig. 14: Geographical distribution and relative abundance of amphipods by RMT 8 samples in November/December 1977.




Fig. 15: Geographical distribution and relative abundance of amphipods by RMT 8 samples in January/February 1976.




Fig. 16: Geographical distribution and relative abundance of amphipods by RMT 8 samples in January/February 1978.




Fig. 17: Geographical distribution and relative abundance of amphipods by RMT 8 samples in January/February 1981.




Fig. 18: Geographical distribution and relative abundance of amphipods by RMT 8 samples in March 1978.




Fig. 19: Geographical distribution and relative abundance of amphipods by RMT 8 samples in March 1981.




Fig. 20: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in November/December 1977.




Fig. 21: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in January/February 1976.






Fig. 22: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in January/February 1978.




Fig. 23: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in January/February 1981.




Fig. 24: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in March 1978.




Fig. 25: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in March 1981.




Fig. 26: Geographical distribution and relative abundance of salps by RMT 8 samples in November/December 1977.




Fig. 27: Geographical distribution and relative abundance of salps by RMT 8 samples in January/February 1976.




Fig. 28: Geographical distribution and relative abundance of salps by RMT 8 samples in January/February 1978.




Fig. 29: Geographical distribution and relative abundance of salps by RMT 8 samples in January/February 1981.




Fig. 30: Geographical distribution and relative abundance of salps by RMT 8 samples in March 1978.




Fig. 31: Geographical distribution and relative abundance of salps by RMT 8 samples in March 1981.

Central Scotia Sea




Fig. 32: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in November/December 1975 (modified from Pommeranz 1978).




Fig. 33: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in November/December 1977 (modified from Wörner 1979).




Fig. 34: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples from January to March 1978 (modified from Wörner 1979).




Fig. 35: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in February 1981 (modified from Nast 1982b).




Fig. 36: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in February/March 1976 (modified from Pommeranz 1978).




Fig. 37: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in November/December 1975.






Fig. 38: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in November/December 1977.




Fig. 39: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples from January to March 1978.




Fig. 40: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in February 1981.




Fig. 41: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in February/March 1976.




Fig. 42: Geographical distribution and relative abundance of amphipods by RMT 8 samples in November/December 1975.




Fig. 43: Geographical distribution and relative abundance of amphipods by RMT 8 samples in November/December 1977.




Fig. 44: Geographical distribution and relative abundance of amphipods by RMT 8 samples from January to March 1978.




Fig. 45: Geographical distribution and relative abundance of amphipods by RMT 8 samples in February 1981.




Fig. 46: Geographical distribution and relative abundance of amphipods by RMT 8 samples in February/March 1976.




Fig. 47: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in November/December 1975.




Fig. 48: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in November/December 1977.




Fig. 49: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples from January to March 1978.




Fig. 50: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in February 1981.




Fig. 51: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in February/March 1976.




Fig. 52: Geographical distribution and relative abundance of salps by RMT 8 samples in November/December 1975.




Fig. 53: Geographical distribution and relative abundance of salps by RMT 8 samples in November/December 1977.






Fig. 54: Geographical distribution and relative abundance of salps by RMT 8 samples from January to March 1978.




Fig. 55: Geographical distribution and relative abundance of salps by RMT 8 samples in February 1981.




Fig. 56: Geographical distribution and relative abundance of salps by RMT 8 samples in February/March 1976.

South Georgia




Fig. 57: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in December 1975 (modified from Pommeranz 1978).




Fig. 58: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in November/December 1977 (modified from Wörner 1979).




Fig. 59: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in January 1978 (modified from Wörner 1979).




Fig. 60: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in March/April 1976 (modified from Pommeranz 1978).




Fig. 61: Geographical distribution and relative abundance of *Euphausia superba* by RMT 8 samples in March/April 1978 (modified from Wörner 1979).




Fig. 62: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in December 1975.




Fig. 63: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in November/December 1977.




Fig. 64: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in January 1978.




Fig. 65: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in March/April 1976.




Fig. 66: Geographical distribution and relative abundance of euphausiids (excl. *Euphausia superba*) by RMT 8 samples in March/April 1978.




Fig. 67: Geographical distribution and relative abundance of amphipods by RMT 8 samples in December 1975.




Fig. 68: Geographical distribution and relative abundance of amphipods by RMT 8 samples in November/December 1977.






Fig. 69: Geographical distribution and relative abundance of amphipods by RMT 8 samples in January 1978.




Fig. 70: Geographical distribution and relative abundance of amphipods by RMT 8 samples in March/April 1976.




Fig. 71: Geographical distribution and relative abundance of amphipods by RMT 8 samples in March/April 1978.




Fig. 72: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in December 1975.




Fig. 73: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in November/December 1977.




Fig. 74: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in January 1978.




Fig. 75: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in March/April 1976.




Fig. 76: Geographical distribution and relative abundance of chaetognaths by RMT 8 samples in March/April 1978.




Fig. 77: Geographical distribution and relative abundance of salps by RMT 8 samples in December 1975.




Fig. 78: Geographical distribution and relative abundance of salps by RMT 8 samples in November/December 1977.




Fig. 79: Geographical distribution and relative abundance of salps by RMT 8 samples in January 1978.




Fig. 80: Geographical distribution and relative abundance of salps by RMT 8 samples in March/April 1976.




Fig. 81: Geographical distribution and relative abundance of salps by RMT 8 samples in March/April 1978.

**Folgende Hefte der Reihe „Berichte zur Polarforschung“ sind bisher erschienen:**

Verkaufspreis/DM

- \* **Sonderheft Nr. 1/1981** – „Die Antarktis und ihr Lebensraum“  
Eine Einführung für Besucher – Herausgegeben im Auftrag von SCAR
- Heft Nr. 1/1982** – „Die Filchner-Schelfeis-Expedition 1980/1981“  
zusammengestellt von Heinz Kohnen 11,50
- Heft Nr. 2/1982** – „Deutsche Antarktis-Expedition 1980/1981 mit FS ‚Meteor‘“  
First International BIOMASS Experiment (FIBEX) – Liste der Zooplankton- und Mikronektonnetzfüge  
zusammengestellt von Norbert Klages 10,—
- Heft Nr. 3/1982** – „Digitale und analoge Krill-Echolot-Rohdatenerfassung an Bord des Forschungs-  
schiffes ‚Meteor‘“ (im Rahmen von FIBEX 1980/81, Fahrtabschnitt ANT III), von Bodo Morgenstern 19,50
- Heft Nr. 4/1982** – „Filchner-Schelfeis-Expedition 1980/81“  
Liste der Planktonfänge und Lichtstärkemessungen  
zusammengestellt von Gerd Hubold und H. Eberhard Drescher 12,50
- \* **Heft Nr. 5/1982** – „Joint Biological Expedition on RRS ‚John Biscoe‘, February 1982“  
by G. Hempel and R. B. Heywood
- \* **Heft Nr. 6/1982** – „Antarktis-Expedition 1981/1982 (Unternehmen ‚Eiswarte‘)“  
zusammengestellt von Gode Gravenhorst
- Heft Nr. 7/1982** – „Marin-Biologisches Begleitprogramm zur Standorterkundung 1979/80 mit MS ‚Polar-  
sirkel‘ (Pre-Site Survey)“ – Stationslisten der Mikronekton- und Zooplanktonfänge sowie der Bodenfischerei  
zusammengestellt von R. Schneppenheim 13,—
- Heft Nr. 8/1983** – „The Post-Fibex Data Interpretation Workshop“  
by D. L. Cram and J.-C. Freytag with the collaboration of J. W. Schmidt, M. Mall, R. Kresse, T. Schwinghammer 10,—
- Heft Nr. 9/1983** – „Distribution of some groups of zooplankton in the inner Weddell Sea in summer 1979/80“  
by I. Hempel, G. Hubold, B. Kaczmaruk, R. Keller, R. Weigmann-Haass 15,—
- Heft Nr. 10/1983** – „Fluor im antarktischen Ökosystem“ – DFG-Symposium November 1982  
zusammengestellt von Dieter Adelung 23,—
- Heft Nr. 11/1983** – „Joint Biological Expedition on RRS ‚John Biscoe‘, February 1982 (II)“  
Data of micronekton and zooplankton hauls, by Uwe Piatkowski 16,—
- Heft Nr. 12/1983** – „Das biologische Programm der ANTARKTIS-I-Expedition 1983 mit FS ‚Polarstern‘“  
Stationslisten der Plankton-, Benthos- und Grundschnepnetzfüge und Liste der Probenahme an Robben  
und Vögeln, von H. E. Drescher, G. Hubold, U. Piatkowski, J. Plötz und J. Voß 14,—
- \* **Heft Nr. 13/1983** – „Die Antarktis-Expedition von MS ‚Polarbjörn‘ 1982/83“ (Sommercampagne zur  
Atka-Bucht und zu den Kraul-Bergen), zusammengestellt von Heinz Kohnen
- \* **Sonderheft Nr. 2/1983** – „Die erste Antarktis-Expedition von FS ‚Polarstern‘ (Kapstadt, 20. Januar 1983 –  
Rio de Janeiro, 25. März 1983)“, Bericht des Fahrleiters Prof. Dr. Gotthilf Hempel
- \* **Sonderheft Nr. 3/1983** – „Sicherheit und Überleben bei Polarexpeditionen“  
zusammengestellt von Heinz Kohnen
- Heft Nr. 14/1983** – „Die erste Antarktis-Expedition (ANTARKTIS I) von FS ‚Polarstern‘ 1982/83“  
(In Vorbereitung)
- Sonderheft Nr. 4/1983** – „On the Biology of Krill *Euphausia superba*“ – Proceedings of the Seminar  
and Report of the Krill Ecology Group, Bremerhaven 12.–16. May 1983, edited by S. B. Schnack 75,—
- Heft Nr. 15/1983** – „German Antarctic Expedition 1980/81 with FRV ‚Walther Herwig‘ and RV ‚Meteor‘“ –  
First International BIOMASS Experiment (FIBEX) – Data of micronekton and zooplankton hauls  
by Uwe Piatkowski and Norbert Klages 22,50
- Sonderheft Nr. 5/1984** – „The observatories of the Georg-von-Neumayer Station“, by Ernst Augstein 8,—
- Heft Nr. 16/1984** – „FIBEX cruise zooplankton data“  
by U. Piatkowski, I. Hempel and S. Rakusa-Suszczewski 19,—
- Heft Nr. 17/1984** – „Fahrtbericht (cruise report) der ‚Polarstern‘-Reise ARKTIS I, 1983“  
von E. Augstein, G. Hempel und J. Thiede 29,—
- Heft Nr. 18/1984** – „Die Expedition ANTARKTIS II mit FS ‚Polarstern‘ 1983/84“,  
Bericht von den Fahrtabschnitten 1, 2 und 3, herausgegeben von D. Fütterer 25,—
- Heft Nr. 19/1984** – „Die Expedition ANTARKTIS II, mit FS ‚Polarstern‘ 1983/84“,  
Bericht vom Fahrtabschnitt 4, Punta Arenas–Kapstadt (ANT-II/4), herausgegeben von H. Kohnen 41,—
- Heft Nr. 20/1984** – „Die Expedition ARKTIS II des FS ‚Polarstern‘ 1984, mit Beiträgen des FS ‚Valdivia‘  
und des Forschungsflugzeuges ‚Falcon 20‘ zum Marginal Ice Zone Experiment 1984 (MIZEX)“  
von E. Augstein, G. Hempel, J. Schwarz, J. Thiede und W. Weigel 42,—