



ORIGINAL RESEARCH

Ontogenetic resource utilization and migration reconstruction with $\delta^{13}\text{C}$ values of essential amino acids in the *Cynoscion acoupa* otolith

Kim Vane¹  | Thomas Larsen²  | Barbara M. Scholz-Böttcher³ | Bernd Kopke³ | Werner Ekau¹¹Leibniz Centre for Tropical Marine Research, Bremen, Germany²Leibniz-Laboratory for Radiometric Dating and Stable Isotope Research, Christian-Albrechts Universität zu Kiel, Kiel, Germany³Institute for Chemistry and Biology of the Marine Environment, University of Oldenburg, Oldenburg, Germany**Correspondence**Kim Vane, Leibniz Centre for Tropical Marine Research, Bremen, Germany.
Email: kim.vane@leibniz-zmt.de**Present address**Kim Vane, Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany.
Email: kim.vane@awi.de or kim.vane@ymail.com**Abstract**

With the increasing anthropogenic impacts on fish habitats, it has become more important to understand which primary resources sustain fish populations. This resource utilization can differ between fish life stages, and individuals can migrate between habitats in search of resources. Such lifetime information is difficult to obtain due to the large spatial and temporal scales of fish behavior. The otolith organic matrix has the potential to indicate this resource utilization and migration with $\delta^{13}\text{C}$ values of essential amino acids (EAAs), which are a direct indication of the primary producers. In a proof-of-concept study, we selected the Acoupa weakfish, *Cynoscion acoupa*, as a model fish species with distinct ontogenetic migration patterns. While it inhabits the Brazilian mangrove estuaries during juvenile stages, it moves to the coastal shelf as an adult. Thus, we expected that lifetime resource utilization and migration would be reflected in $\delta^{13}\text{C}_{\text{EAA}}$ patterns and baseline values in *C. acoupa* otoliths. By analyzing the *C. acoupa* otolith edges across a size range of 12–119 cm, we found that baseline $\delta^{13}\text{C}_{\text{EAA}}$ values increased with size, which indicated an estuarine to coastal shelf distribution. This trend is highly correlated with inorganic $\delta^{13}\text{C}$ values. The $\delta^{13}\text{C}_{\text{EAA}}$ patterns showed that estuarine algae rather than mangrove-derived resources supported the juvenile *C. acoupa* populations. Around the juvenile size of 40 cm, resource utilization overlapped with those of adults and mean baseline $\delta^{13}\text{C}_{\text{EAA}}$ values increased. This trend was confirmed by comparing otolith core and edges, although with some individuals potentially migrating over longer distances than others. Hence, $\delta^{13}\text{C}_{\text{EAA}}$ patterns and baseline values in otoliths have great potential to reconstruct ontogenetic shifts in resource use and habitats. The insight could aid in predictions on how environmental changes affect fish populations by identifying the controlling factors at the base of the food web.

KEYWORDS

Amazon, carbon isotopes, life history, organic matrix, otolith chemistry

1 | INTRODUCTION

Fish populations are increasingly influenced by human activities such as removal of estuarine vegetation for rural development, discharge of chemicals, climate change, and ocean acidification (Cheung, Brodeur, Okey, & Pauly, 2015; Halpern et al., 2008; Jackson, 2001; Nagelkerken, Russell, Gillanders, & Connell, 2015). To understand the impact of these habitat changes to the stability of fish populations, it is imperative to know which resources sustain them. By providing the resources at the bottom of the food web, primary producers are a major factor in determining the productivity at lower trophic levels and thus also predatory fish populations at higher trophic levels (Lynam et al., 2017; Pauly & Christensen, 1995). The use of resources can vary during fish life stages as is indicated by their utilization of distinct habitats or niches in an ecosystem (Huijbers, Nagelkerken, Debrot, & Jongejans, 2013; Kimirei et al., 2015). Moreover, fish actively undertake periodic migrations to synchronize their spawning time toward seasons with optimal resources for themselves and their offspring (Dingle & Drake, 2007; Nakazawa & Yamamura, 2006). Therefore, each life stage is sensitive to changes among primary producers, due to, for example, anthropogenic changes in the ecosystem, which affect the entire population. Shifts among primary producers affecting fish populations are not well characterized, as observational studies are often inconclusive in identifying key primary producers or habitat value for a fish species. Detecting a fish in a particular habitat does not directly identify the primary producers sustaining the food web it feeds on (Kruitwagen, Nagelkerken, Lugendo, Mgamma, & Bonga, 2010; Seitz, Wennhage, Bergstrom, Lipcius, & Ysebaert, 2014). This lack of knowledge also impairs effective conservation strategies for maintaining fish populations (Sheaves, Baker, Nagelkerken, & Connolly, 2015).

Characterizing resource utilization by juvenile stages is key for understanding population dynamics as they determine the recruitment of the adult population. How habitats are used by juveniles has been difficult to resolve and can be attributed to the complications of tracking especially small juveniles between habitats, which can occur periodically and over large spatial scales (Beck et al., 2001; Dahlgren et al., 2006; Gillanders, Able, Brown, Eggleston, & Sheridan, 2003). For this reason, fish otolith inorganic chemistry has been often utilized to qualify the movements of fish between habitats as otoliths provide a complete record of the fish's lifetime (Campana, 1999; Elsdon et al., 2008). The otolith calcium carbonate is accreted continuously from the early embryonic stage throughout the individual's life with often clear growth rings that are an indication of its age (Campana & Neilson, 1985). Simultaneously, the otolith calcium carbonate matrix is metabolically inert and retains the environmental chemistry in terms of minor and trace elements and their isotopes (e.g., strontium, barium, bulk $\delta^{13}\text{C}$, bulk $\delta^{18}\text{O}$). By interpreting elemental concentrations or isotopic values in the otolith, one could deduce whether the fish individual traversed distinct chemical environments such as freshwater and marine water habitats (Bath et al., 2000; Secor, Henderson-Arzapalo, & Piccoli, 1995). However, there is growing evidence that the physiological

and metabolic pathways of fish have implications for the elemental concentrations and bulk $\delta^{13}\text{C}$ values in the otolith (Campana, 1999; Geffen, 2012; Radtke, Williams, & Hurley, 1987; Solomon et al., 2006; Sturrock et al., 2014). This causes uncertainty in the extent to which otolith inorganic chemistry can be interpreted for movement patterns and what is caused by physiological and metabolic changes (Chang & Geffen, 2013; Grønkvær, 2016).

The recent utilization of the otolith organic part, consisting primarily of proteins, could provide a more direct and unperturbed link between fish and its ambient environment (McMahon, Fogel, et al., 2011). Amino acids (AAs) are the building blocks of proteins, and about half of the 20 protein AAs are essential for metazoans, because they cannot synthesize them de novo (Howland et al., 2003). Therefore, metazoans depend on the essential amino acids (EAAs) through dietary sources. These EAAs can be traced to their biosynthetic origins such as algae, bacteria, fungi, and vascular plants. The carbon isotopic compositions or $\delta^{13}\text{C}$ baseline values of EAAs, among primary producers, display geographical variations (i.e., isoscapes) and can therefore indicate fish movement and habitat connectivity (McMahon, Berumen, & Thorrold, 2012; Vokhshoori, Larsen, & McCarthy, 2014). Yet, each primary producer can have distinct biosynthetic pathways that lead to characteristic isotopic fractionations or $\delta^{13}\text{C}$ patterns of individual EAAs ($\delta^{13}\text{C}_{\text{EAA}}$) that can be used to identify the primary producer. These source diagnostic $\delta^{13}\text{C}$ patterns or fingerprints of EAAs can be used as a marker of primary producers regardless of the $\delta^{13}\text{C}$ baseline variations (Arthur, Kelez, Larsen, Choy, & Popp, 2014; Larsen et al., 2013). Thus, the otolith temporal increments and incorporated EAAs constitute an archive of lifetime resource utilization and migration through the individual fish life stages. However, only a limited amount of studies have looked into the potential of using the otolith organic matrix (Grønkvær et al., 2013; McMahon, Berumen, Mateo, Elsdon, & Thorrold, 2011; McMahon, Fogel, et al., 2011; McMahon et al., 2012).

The known distribution of the Acoupa weakfish, *Cynoscion acoupa*, from mangrove estuaries in the early life stages to offshore coastal shelf areas as adults (Barletta, Barletta-Bergan, & Saint-Paul, 1998; Barletta, Barletta-Bergan, Saint-Paul, & Hubold, 2003) suggests an ontogenetic migration between the two habitats and changes in resource utilization. Although it is a commercially important species occurring along the entire coast of Brazil, little is known about its foraging ecology or migration. Thus, $\delta^{13}\text{C}_{\text{EAA}}$ baseline values and patterns from *C. acoupa* otoliths could provide more insight into its ontogenetic resource utilization and migration. We collected fish otoliths of *C. acoupa* individuals across a size range to identify habitat change and primary resource utilization along the ontogenetic development of this species. The main aim of this study was therefore to investigate the potential of $\delta^{13}\text{C}_{\text{EAA}}$ analysis of otoliths to extract lifetime records of movement and resource utilization of *C. acoupa*. We tested whether baseline $\delta^{13}\text{C}_{\text{EAA}}$ values in otolith edges of *C. acoupa* individuals at different lengths are consistent with the isotopic trend displayed by otolith inorganic $\delta^{13}\text{C}$ values. We posit that $\delta^{13}\text{C}_{\text{EAA}}$ patterns in combination with baseline $\delta^{13}\text{C}_{\text{EAA}}$ values will reveal *C. acoupa* lifetime resource utilization and

distribution from predominantly estuarine resources during juvenile stages to coastal shelf resources during mature stages. To test this, we analyzed otolith edges among different size classes and compared individual otolith cores and edges.

2 | MATERIALS AND METHODS

2.1 | Species and sampling location

The Acoupa weakfish, or *C. acoupa*, is a demersal fish that is caught by an artisanal fishery up to ~125 cm total length and weigh ~16 kg where adults are exclusively caught on the coastal shelf (Barletta et al., 1998; de Matos & Lucena, 2017). This sciaenid species occurs on the coastal shelf along the entire coast of Brazil and utilizes the mangrove estuaries as larvae and juveniles (Barletta et al., 2003; Barletta-Bergan, Barletta, & Saint-Paul, 2002a; Lima, Barletta, & Costa, 2015, Figure 1). In northern Brazil, *C. acoupa* spawns offshore at the onset and during the wet seasons and sexual maturation occurs ~40 cm (Almeida, Santos, Sousa, Carvalho Neta, & Andrade, 2016). It has a carnivorous diet, which consists mainly of shrimp and fish as well as small crustaceans and polychaetes in the juvenile stage (Ferreira et al., 2016).

Sampling of *C. acoupa* occurred only in Pará, northern Brazil, which receives a high annual precipitation of over 2,000 mm and leads to high freshwater discharges by the Amazon River. Consequently, in the wet season a high-nutrient load is dispersed several kilometers offshore in a northward direction over a wide coastal shelf (Bustamante et al., 2015; Knoppers, Ekau, & Figueiredo, 1999; Smith & Demaster, 1996). The estuarine environment harbors extensive mangrove deltas and is strongly macrotidal with amplitudes of ~4 m (Schaeffer-Novelli et al., 1990).

2.2 | Sampling locations

Sampling of *C. acoupa* otoliths was conducted in Bragança, one of the main landing ports in the state of Pará, northern Brazil. Twenty juvenile *C. acoupa* individuals of 11–40 cm standard length (SL) were collected from local fishermen at the Caeté River, and 32 adults of 56–119 cm SL were sampled at the fish market in Bragança. Otoliths were removed, cleaned with demi water, and stored dry. Primary producers were collected at the Caeté River mouth close to where the juvenile fish were collected by the fishermen. No seagrass or coral reefs are found in this region. Leaves of the *Rhizophora mangle* mangrove tree, a rhodophyte growing on the mangrove roots *Bostrychia* sp., the freshwater algae *Spirogyra* sp., and degraded brown mangrove leaves, representing the bacterial community, were collected in the Caeté estuary. Coastal zooplankton, consisting predominantly of *Acartia tonsa*, was collected with a plankton net (mesh size 300 μ m) to represent the coastal phytoplankton community. All primary producer samples were rinsed with MilliQ water, freeze-dried, and stored in a -20° freezer.

2.3 | Otolith preparation

Juvenile otoliths of *C. acoupa* individuals between 12 and 25 cm SL were used whole, cleaned in an ultrasonic bath with ultrapure water, and dried in the oven for 3 hr at 40°C. Subsequently, otoliths were homogenized with mortar and pestle and stored in a combusted glass vial. Otoliths of juvenile and adults between 25 and 119 cm were embedded in Araldite 2020 epoxy resin. Embedded otoliths were cut with a diamond saw through the center for a section of approximately 2 mm thickness. Sections were then glued with epoxy resin on a glass slide. Apart from an acetone wipe, the section surface was untreated to prevent contamination. The edges of these sectioned otoliths were micromilled for a 5 mg sample. To gain insight into resource utilization in the early life stage and latest life stage of seven individuals between 38 and 110 cm SL, the cores were micromilled alongside the otolith edges. Cores were micromilled according to the dimensions of the smallest otolith in the collection originating from a 12 cm SL *C. acoupa* individual.

2.4 | Otolith bulk $\delta^{13}\text{C}$ analysis

The inorganic calcium carbonate $\delta^{13}\text{C}$ values from 42 homogenized *C. acoupa* otoliths and micromilled otolith edges were measured on a Finnigan MAT 251 gas isotope ratio mass spectrometer. This was connected to a Kiel III automated carbonate preparation device where samples react with phosphoric acid at 74°C. Data are reported in delta notation versus Vienna Pee Dee Belemnite standard (V-PDB). The instrument was calibrated against the house standard (ground Solnhofen limestone), which in turn was calibrated against the NBS 19 standard reference material. Over the measurement period, the standard deviations of the house standard were 0.04‰ for $\delta^{13}\text{C}$ values.

2.5 | Otolith $\delta^{13}\text{C}$ -AA analysis

The 5 mg of all homogenized and micromilled otolith powder and 1 mg primary producer samples were hydrolyzed with 100 μ l of 6M HCl per mg of sample in a microreaction vessel and 4 ml glass vial, respectively, with a reference of 20 μ l 6-aminocaproic acid at the same time. After flushing the vials with nitrogen, the samples were heated at 110°C for 20 hr. The acid was then evaporated in a heating block at 110°C for approximately 30 min. After acid hydrolysis, only the primary producer samples were purified with a DOWEX 50WX8 (hydrogen form, 100–200 mesh) column and eluted with 2M ammonium hydroxide. This was then evaporated to dryness in a 110°C aluminum block for approximately 1 hr. The samples were then redissolved in 200 μ l of 0.1 M HCl, and 100 μ l was transferred to a micro-reaction vessel. All otolith and primary producer samples were then derivatized with the methoxycarbonyl esterification protocol (Walsh, He, & Yarnes, 2014). This included the addition of 35 μ l methanol, 30 μ l pyridine, and 15 μ l methyl chloroformate. With a Hamilton syringe, 100 μ l of chloroform was added, mixed, and rested for 5 min to separate into organic layers. The bottom chloroform



FIGURE 1 A juvenile *Cynoscion acoupa* of 32 cm total length from the Caeté River in Pará, Brazil

layer was removed and transferred to a GC vial with 250 μ l insert with a small amount of sodium sulfate. To prevent the sodium sulfate grains getting into the injector of the GC-c-IRMS, 40 μ l was again transferred to a new GC vial with 250 μ l insert and 4 μ l of $C_{12:0}/C_{20:0}$ FAME standard was added.

A 2 μ l aliquot of the sample was then injected in an Agilent 6890 GC with a VF-WAXms column (30 m, 0.25 μ m i.d., 0.25 μ m film thickness) and interfaced to a MAT253 IRMS (Finnigan MAT) via a GC-C II combustion interface. The GC temperature program utilized was 60°C (2 min), increased with 3°C/s to 245°C, and held for 20.3 min. All samples were injected via a cold injection system (CIS 4; Gerstel) in the splitless mode.

2.6 | Data analysis

We obtained consistently good chromatography with sufficient chromatographic baseline separation across all samples for five EAAs, valine (Val), isoleucine (Iso), threonine (Thr), phenylalanine (Phe), and lysine (Lys). The measured $\delta^{13}C_{EAA}$ values were corrected for kinetic fractionation during derivatization according to Docherty, Jones, and Evershed (2001). We followed a dual approach to present and evaluate the $\delta^{13}C_{EAA}$ data; one based on baseline $\delta^{13}C_{EAA}$ values (the corrected $\delta^{13}C_{EAA}$ values) and the other on mean normalized $\delta^{13}C_{EAA}$ values. The mean normalized values are calculated by subtracting each individual $\delta^{13}C_{EAA}$ value from the mean $\delta^{13}C$ values of all EAAs for each sample. While baseline $\delta^{13}C_{EAA}$ values reflect spatial-temporal isotopic variation in the environment, normalized $\delta^{13}C_{EAA}$ values track the biosynthetic origins of EAAs to algae, bacteria, fungi, and vascular plants independently of environmental conditions (Larsen et al. 2015). These source diagnostic $\delta^{13}C_{EAA}$ patterns are also denoted as $\delta^{13}C_{EAA}$ fingerprints as in Larsen, Taylor, Leigh, and O'Brien (2009); Larsen

et al. (2013). To identify these EAA sources, we compared $\delta^{13}C_{EAA}$ patterns between *C. acoupa* otoliths of individuals at different life stages and collected primary producers by performing multivariate analyses in R version 3.4.3 (R-Development-Core-Team 2017). For assessing dietary contribution of primary producer sources to *C. acoupa* individuals, we applied principal component analysis (PCA; R: vegan) to the normalized $\delta^{13}C_{EAA}$ values from otoliths and primary producers. We performed covariance matrix PCA to preserve variance as the range and scale of variables are in the same units of measure. We also created a classification model, linear discriminant function analysis (LDA; R: MASS), to determine the probability of dietary contributions to consumers. It is not necessary to mean normalize $\delta^{13}C_{EAA}$ values prior to applying LDA, because the LDA procedure that scales observations to discriminant functions already normalizes them to ensure that within groups covariance matrix is spherical (Venables & Ripley, 2002). LDA is also useful for selecting the variables that are the most informative for discriminating between groups or classes. Similarly, univariate analysis performed on the output from multivariate analysis of variance (MANOVA; R: manova) can also be applied to test which dependent variables ($\delta^{13}C_{EAA}$ values) are significantly different between groups (R: summary.aov). In conjunction with MANOVA, we used Pillai's trace to test the null hypothesis that groups have a common centroid in a dependent variable vector space. A rejection of this hypothesis entails that the groups have significantly different $\delta^{13}C_{EAA}$ patterns or baseline values. To compare bulk inorganic $\delta^{13}C$ values and mean baseline $\delta^{13}C_{EAA}$ values, Spearman's correlation was applied due to non-normally distributed data.

3 | RESULTS

3.1 | Otolith bulk $\delta^{13}C$ versus mean $\delta^{13}C_{EAA}$ values

Inorganic $\delta^{13}C$ values and mean baseline $\delta^{13}C_{EAA}$ values in the same otoliths of *C. acoupa* individuals ranging between 12 and 119 cm SL showed the same trend of becoming more positive with increasing SL. These $\delta^{13}C$ values were tightly correlated ($\rho = 0.70$, $p < 0.001$, Figure 2a), but with the inorganic fraction being 10‰ more enriched than the organic EAA fraction. Moreover, the range in mean baseline $\delta^{13}C_{EAA}$ values was greater than that in bulk inorganic $\delta^{13}C$ values (4‰ vs. 2.5‰ respectively, Figure 2b).

3.2 | $\delta^{13}C_{EAA}$ patterns in otoliths and primary producers

Coastal primary producers as mangrove root rhodophytes and degraded and fresh mangrove leaves had overall more negative baseline $\delta^{13}C_{EAA}$ values than the aquatic freshwater algae and phytoplankton (Figure 3a). Also, baseline $\delta^{13}C_{EAA}$ values in otoliths from 12 to 119 cm SL individuals displayed an overall increasing trend from small to large individuals (Figure 3b) with juvenile (<40 cm) and adult *C. acoupa* groups (>55 cm) being significantly different (Pillai's trace = 0.67, $F_{5,47} = 18.9$; $p < 0.001$). For all EAAs,




FIGURE 2 (a) Mean baseline $\delta^{13}\text{C}_{\text{EAA}}$ values versus the inorganic $\delta^{13}\text{C}$ values measured in the same otolith material ($N = 42$). (b) Mean otolith baseline $\delta^{13}\text{C}_{\text{EAA}}$ values ($N = 52$) and a subsample of the same otoliths measured as inorganic $\delta^{13}\text{C}$ values ($N = 42$) from individuals with a standard length from 12 to 119 cm

adult otolith edges were more ^{13}C -enriched than juvenile otoliths ($p < 0.05$).

We then investigated the extent in which baseline independent $\delta^{13}\text{C}_{\text{EAA}}$ patterns can differentiate among otoliths of adults and juveniles and the various primary producer groups. The primary producers display particularly contrasting $\delta^{13}\text{C}_{\text{EAA}}$ patterns in regard to Val, Phe, and Thr (Figure 3c). In comparison, $\delta^{13}\text{C}_{\text{EAA}}$ patterns were less distinct between the juvenile and adult *C. acoupa* groups (Figure 3d). In the PCA (Figure 4a), Val, Phe, and Thr were strongly correlated with the first principal component separating freshwater from marine resources. To test whether juvenile and adult *C. acoupa* have different $\delta^{13}\text{C}_{\text{EAA}}$ patterns, we followed a two-step approach. We first identified and removed the EAAs that do not have significantly different mean normalized $\delta^{13}\text{C}$ values between juvenile and adult otoliths, and then performed a MANOVA test with the remaining EAAs. After removing Ile and Val, the least informative variables for discriminating between adults and juveniles, we found that the juvenile and adult *C. acoupa* groups were significantly different (Pillai's trace = 0.55, $F_{3,49} = 19.5$; $p < 0.001$). According to the PCA (Figure 4a), Val and Ile were also the least correlated with the second principal component separating algal- and mangrove-derived resources.

The PCA results showed that $\delta^{13}\text{C}_{\text{EAA}}$ patterns of fish otoliths resembled coastal phytoplankton and freshwater algae more than mangrove root rhodophytes and degraded and fresh mangrove leaves. To visualize the distributional and resource utilization information that can be obtained from both mean baseline $\delta^{13}\text{C}_{\text{EAA}}$ values and $\delta^{13}\text{C}_{\text{EAA}}$ patterns, respectively, we constructed a PCA that incorporates both indicators. In the PCA based on $\delta^{13}\text{C}_{\text{EAA}}$ patterns of adult and juvenile otoliths and the primary producers (Figure 4a), we color-coded the principal component points according to their mean baseline $\delta^{13}\text{C}_{\text{EAA}}$ values (Figure 4b). Red colors represent more positive baseline $\delta^{13}\text{C}_{\text{EAA}}$ values and vice versa for blue colors (Figure 4b). The color gradient showed that

adults have more positive baseline $\delta^{13}\text{C}_{\text{EAA}}$ values than juveniles and coastal resources and that about half of the juveniles share baseline $\delta^{13}\text{C}_{\text{EAA}}$ values with coastal resources (Figure 4b).

3.3 | $\delta^{13}\text{C}_{\text{EAA}}$ values in otolith core and edge

The thick sectioning of northern *C. acoupa* otoliths did not allow us to detect incremental structures. Instead, we micromilled cores according to the otolith size dimensions of a 12 cm SL individual to get an insight into early and later life stages of seven individuals with 38–110 cm SL. Consistently lower baseline $\delta^{13}\text{C}_{\text{EAA}}$ values were observed in the individual otolith cores than in the edges (ANOVA, $p < 0.01$ for Ile, Lys, Thr, Val; $p < 0.05$ for Phe; Figure 5a). We found that only Phe and Thr were significantly different between the two groups (ANOVA, $p < 0.01$; Figure 5b). This finding is consistent with Phe and Thr being among the most informative EAAs for distinguishing between juvenile and adult *C. acoupa* (Figure 4a). However, in a dependent variable vector space based on all five EAAs, it was indicated that core and edge otoliths had significantly different $\delta^{13}\text{C}_{\text{EAA}}$ patterns (Pillai's trace = 0.97, $F_{5,8} = 60.5$; $p < 0.001$).

To investigate individual lifetime use of estuarine and coastal shelf resources, we compared baseline $\delta^{13}\text{C}_{\text{EAA}}$ values between core and edge otolith samples. For the training data, we selected five juveniles and five adults with the lowest and highest mean baseline $\delta^{13}\text{C}_{\text{EAA}}$ values, respectively. As for the variables, we used the three most informative EAAs (Lys, Phe, Thr) for distinguishing between estuarine and coastal shelf resources. This classification model was applied to predict ontogenetic habitat shifts of seven *C. acoupa* individuals with different SL (Figure 6). Practically all individuals, excluding SL38, have core $\delta^{13}\text{C}_{\text{EAA}}$ values that tended more toward estuarine resources than the edge measurements. However, some core or juvenile values, SL71 and SL87, overall inclined more




FIGURE 3 Individual baseline $\delta^{13}\text{C}$ values of Val, Ile, Thr, Phe, and Lys with standard deviation from (a) primary producers and (b) whole fish otoliths from *Cynoscion acoupa* individuals with 12–25 cm SL and otolith edges from individuals with 25–119 cm SL. All measurements are based on triplicate injections. The mean normalized $\delta^{13}\text{C}_{\text{EAA}}$ values for (c) primary producers and (d) *C. acoupa* fish otoliths

toward coastal shelf baseline $\delta^{13}\text{C}_{\text{EAA}}$ values and approach edge or adult values (Figure 6).

4 | DISCUSSION

We obtained evidence of ontogenetic estuarine to coastal shelf migration by *C. acoupa* individuals from both inorganic $\delta^{13}\text{C}$ and

mean baseline $\delta^{13}\text{C}_{\text{EAA}}$ values of otoliths. Yet, among individuals in juvenile and adult stages, considerable variations in both $\delta^{13}\text{C}$ tracers were observed. Variations in $\delta^{13}\text{C}$ value are not unexpected in tissues of a species residing in the highly variable Amazon region across a large distribution area with potential to move over considerable distances. In the wet season, the Amazon River disperses large amounts of freshwater into the coastal areas and several kilometers offshore toward the north (Loick-Wilde et al., 2016; Weber




FIGURE 4 (a) Principal component analysis of $\delta^{13}\text{C}_{\text{EAA}}$ patterns in adult ($n = 32$) and juvenile ($n = 20$) otoliths (open symbols) and primary producers (filled symbols). Of the four principal components, the first two account for 88% of the variation (49.7% and 38.3%, respectively). (b) PCA coordinates of $\delta^{13}\text{C}_{\text{EAA}}$ patterns displayed simultaneously with baseline $\delta^{13}\text{C}_{\text{EAA}}$ values as a color gradient with warm colors signifying ^{13}C -enriched samples and vice versa for cool colors

et al., 2017). This can cause substantial spatial differences in $\delta^{13}\text{C}$ baselines (Dittmar, Hertkorn, Kattner, & Lara, 2006) across areas which *C. acoupa* individuals traverse. These geographical, seasonal, and possibly long-term or sudden changes in $\delta^{13}\text{C}$ baseline values that can occur in the Amazon complicate the deduction of migration and resource utilization over long timescales as represented by otolith records. Disentangling metabolic change, diet, and DIC contributions to otolith inorganic carbon has been complicated (Elsdon, Ayvazian, McMahon, & Thorrold, 2010; Nelson, Hanson, Koenig, & Chanton, 2011; Solomon et al., 2006) and impeded the deductions about fish migration from otolith records (Jamieson, Schwarcz, & Bratley, 2004; Kalish, 1991; Schwarcz et al., 1998). Consequently, otolith inorganic $\delta^{13}\text{C}$ trends are poor indicators for identifying $\delta^{13}\text{C}$ baseline or resource changes. Source diagnostic $\delta^{13}\text{C}_{\text{EAA}}$ patterns can overcome these $\delta^{13}\text{C}$ baseline variations because $\delta^{13}\text{C}_{\text{EAA}}$ patterns of primary producers remain largely constant despite the $\delta^{13}\text{C}$ of the environmental conditions or spatial separation of primary producers (Larsen et al., 2009, 2013).

A pattern of lifetime resource utilization and distribution from estuary to coastal shelf by *C. acoupa* with increasing length was

displayed by both $\delta^{13}\text{C}_{\text{EAA}}$ baseline values and patterns. The PCA results depict that the resource utilization of juvenile and adult *C. acoupa* gradually become distinct and overlap at intermediate life stages, which could indicate that there is a regular movement between estuarine and coastal shelf environments by juveniles. An occurrence that is not unlikely in macrotidal Amazon estuaries where tides and body size of several fish species were observed to facilitate movement in and out of mangrove creeks (Brenner & Krumme, 2007; Krumme, Calderón, & Echterhoff, 2014). However, some otolith core measurements tended more toward coastal shelf resources and could imply that not all *C. acoupa* juvenile stages are spent within estuarine mangrove habitats. Estuarine resource use was also indicated by $\delta^{13}\text{C}_{\text{EAA}}$ patterns in some otolith adult stages of *C. acoupa* although adults are not observed within estuaries. We speculate that some adults inhabiting the coastal shelf feed on fish that migrate out of the estuaries onto the coastal shelf. Visual census and a $\delta^{13}\text{C}_{\text{EAA}}$ application also indicated that multiple species, which are believed to make ontogenetic migrations from mangrove or seagrass beds to coral reefs, also spend their early life stages in various habitats or complete their life cycle on




FIGURE 5 (a) A subsample of otoliths from differently sized individuals measured for baseline $\delta^{13}\text{C}_{\text{EAA}}$ values in core (C) and edge (E) of the otoliths. Otolith cores were microdrilled according to the dimensions of an otolith from an individual of 12 cm SL and are shaded in gray. (a) Individual $\delta^{13}\text{C}$ values of Val, Ile, Thr, Phe, and Lys with standard deviation. (b) Calculated mean normalized $\delta^{13}\text{C}_{\text{EAA}}$ values from the otolith core and edges

offshore coral reefs (Kimirei et al., 2011; McMahon et al., 2012). Intrapopulation life history differences can be important for the population's persistence when facing environmental disturbances and can be a result of resource availability, food–predation risk trade-offs, and spatial overlap of food webs (Bolnick et al. 2003; 2011).

Our results also show that coastal phytoplankton and freshwater algae contribute the most as resources to the juvenile *C. acoupa* community and that direct contributions from mangrove-derived resources are negligible. This agrees with




FIGURE 6 Linear discriminant function analysis for predicting use of coastal shelf and estuarine resources based on baseline $\delta^{13}\text{C}$ values of Phe, Lys, and Thr, which are the independent variables displayed on the right as coefficients (crosses). Based on the training data comprising of juvenile and adult *Cynoscion acoupa* individuals, the model predicted that in most cases that EAAs from otolith cores (down-pointing triangles) were more closely associated with estuarine resources than otolith edges (up-pointing triangles), which were more associated with coastal shelf resources. Numbers under the triangles represent standard length in cm for each individual from which the otolith was obtained

the spawning behavior of *C. acoupa* in North Brazil, which occurs with the rainy seasons (Almeida et al., 2016). Large numbers of larval and juvenile *C. acoupa* coincide with large blooms in zooplankton and shrimp populations in the mangrove estuaries, which often rely on planktonic and algal diets (Barletta-Bergan, Barletta, & Saint-Paul, 2002b; Barletta-Bergan et al., 2002a; Ferreira et al., 2016; Lima et al., 2015). Phytoplankton blooms are also enhanced by the increased nutrient concentrations in the Amazon estuaries during the wet season (Dittmar & Lara, 2001; Santos et al., 2008; Smith & Demaster, 1996). However, other studies also indicated that mangrove carbon often does not directly contribute to sustaining the juvenile population (Igulu, Nagelkerken, van der Velde, & Mgaya, 2013; Kruitwagen et al., 2010; Melville & Connolly, 2003). Indirectly, mangroves can support fish populations by providing refuge to juvenile fish and supplying the coastal area with nutrients and mangrove carbon through the microbial and crustacean community (Bouillon et al., 2008; Dittmar et al., 2006; Nordhaus, Wolff, & Diele, 2006; Schories et al., 2003). Carbon from mangrove forests can therefore be an important, albeit indirect, factor in the primary production of coastal and offshore ecosystems supporting *C. acoupa* populations.

To gain a complete insight into the contribution of primary producers to estuarine and offshore fish food webs, all possible primary producers should be collected in the immediate environment and migration area of the species. This would also include the contribution of bacteria and fungi, which have been indicated to play an important role in the diet of first trophic-level consumers (Pascal, Dubois, Goffette, & Lepoint, 2017; Steffan et al., 2017; Yi et al., 2017). Due to time and funding constraints, such samples were not included in the current study, but future studies could take full advantage of the source diagnostic ability of $\delta^{13}\text{C}_{\text{EAA}}$ patterns to distinguish among algal-, bacterial-, fungal-, and plant-derived EAA sources (Larsen et al., 2013). While our results provide a general assessment of *C. acoupa* resource utilization, they concur with previous studies that also found insubstantial direct contributions of mangrove leaves to estuarine fish food webs (Iglu et al., 2013; Kruitwagen et al., 2010; Melville & Connolly, 2003) and the large direct contribution of aquatic algae such as phytoplankton and freshwater algae. This is a strong confirmation that $\delta^{13}\text{C}_{\text{EAA}}$ patterns in otoliths are an indicator of lifetime resource utilization by individual fish.

In summary, $\delta^{13}\text{C}_{\text{EAA}}$ measurements of the otoliths organic matrix display the resource utilization and migration of *C. acoupa* individuals. Baseline $\delta^{13}\text{C}_{\text{EAA}}$ values indicated its known coast to offshore distribution pattern and source diagnostic $\delta^{13}\text{C}_{\text{EAA}}$ patterns displayed a gradual change and overlap in resource utilization of *C. acoupa* juveniles and adults. The otolith organic matrix, hence, has great potential to be used as ontogenetic records with compound specific isotope analysis of $\delta^{13}\text{C}$ in EAAs. As $\delta^{13}\text{C}_{\text{EAA}}$ patterns of primary producers seem unaffected by $\delta^{13}\text{C}$ baseline changes or environmental factors (Larsen et al. 2015), they could be utilized on archeological otoliths to identify changes in resource utilization at different life stages of past fish populations. A temporal comparison based on otolith archives could give us insight into how resource utilization by fish changes in response to habitat degradation and anthropogenic estuarine eutrophication.

ACKNOWLEDGMENTS

Otolith collections in this study were conducted under Sistema de Autorização e Informação em Biodiversidade (SISBIO) field permit number 51486-1 issued by Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) in Brazil.

CONFLICT OF INTEREST

None declared.

AUTHORS' CONTRIBUTIONS

KV conceived the ideas and designed the methodology; KV, BSB, BK, and WE collected the data; KV and TL analyzed the data and led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

DATA ACCESSIBILITY

Data available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.8q7h4g7>

ORCID

Kim Vane  <http://orcid.org/0000-0001-8172-7831>

Thomas Larsen  <http://orcid.org/0000-0002-0311-9707>

REFERENCES

- Almeida, Z. S., Santos, N. B., Sousa, H. L., Carvalho Neta, R. N. F., & Andrade, T. S. O. M. (2016). Biologia reprodutiva da pescada amarela (*Cynoscion acoupa*) capturada na Baía de São Marcos, Maranhão, Brasil. *Biota Amazônia*, 6, 46–54. <https://doi.org/10.18561/2179-5746/biotaamazonia>
- Arthur, K. E., Kelez, S., Larsen, T., Choy, C. A., & Popp, B. N. (2014). Tracing the biosynthetic source of essential amino acids in marine turtles using $\delta^{13}\text{C}$ fingerprints. *Ecology*, 95, 1285–1293. <https://doi.org/10.1890/13-0263.1>
- Barletta, M., Barletta-Bergan, A., & Saint-Paul, U. (1998). Description of the fisheries structure in the mangrove-dominated region of Bragança (state of Para, North Brazil). *Ecotropica*, 4, 41–53.
- Barletta, M., Barletta-Bergan, A., Saint-Paul, U., & Hubold, G. (2003). Seasonal changes in density, biomass, and diversity of estuarine fishes in tidal mangrove creeks of the lower Caeté Estuary (northern Brazilian coast, east Amazon). *Marine Ecology Progress Series*, 256, 217–228. <https://doi.org/10.3354/meps256217>
- Barletta-Bergan, A., Barletta, M., & Saint-Paul, U. (2002a). Structure and seasonal dynamics of larval fish in the Caeté River estuary in North Brazil. *Estuarine, Coastal and Shelf Science*, 54, 193–206. <https://doi.org/10.1006/ecss.2001.0842>
- Barletta-Bergan, A., Barletta, M., & Saint-Paul, U. (2002b). Community structure and temporal variability of ichthyoplankton in North Brazilian mangrove creeks. *Journal of Fish Biology*, 61, 33–51. <https://doi.org/10.1111/j.1095-8649.2002.tb01759.x>
- Bath, G. E., Thorrold, S. R., Jones, C. M., Campana, S. E., McLaren, J. W., & Lam, J. W. H. (2000). Strontium and barium uptake in aragonitic otoliths of marine fish. *Geochimica et Cosmochimica Acta*, 64, 1705–1714. [https://doi.org/10.1016/S0016-7037\(99\)00419-6](https://doi.org/10.1016/S0016-7037(99)00419-6)
- Beck, M. W., Heck, K. L., Able, K. W., Childers, D. L., Eggleston, D. B., Gillanders, B. M., ... Weinstein, M. P. (2001). The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates. *BioScience*, 51, 633–641. [https://doi.org/10.1641/0006-3568\(2001\)051\[0633:TICAMO\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0633:TICAMO]2.0.CO;2)
- Bouillon, S., Borges, A. V., Castañeda-Moya, E., Diele, K., Dittmar, T., Duke, N. C., ... Twilley, R. R. (2008). Mangrove production and carbon sinks: A revision of global budget estimates. *Global Biogeochemical Cycles*, 22, 1–12.
- Bolnick, D. I., Amarasekare, P., Araujo, M. S., Bürger, R., Levine, J. M., Novak, M., Rudolf, V. H. W., Schreiber, S. J., Urban, M. C., & Vasseur, D. A. (2011). Why intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution*, 26, 183–192. <https://doi.org/10.1016/j.tree.2011.01.009>
- Bolnick, D. I., Svanbäck, R., Fordyce, J. A., Yang, L. H., Davis, J. M., Hulseley, C. D., & Forister, M. L. (2003). The ecology of individuals: Incidence and implications of individual specialization. *The American Naturalist*, 161, 1–28. <https://doi.org/10.1086/343878>
- Brenner, M., & Krumme, U. (2007). Tidal migration and patterns in feeding of the four-eyed fish *Anableps anableps* L. in a north Brazilian mangrove. *Journal of Fish Biology*, 70, 406–427. <https://doi.org/10.1111/j.1095-8649.2007.01313.x>

- Bustamante, M. M. C., Martinelli, L. A., Pérez, T., Rasse, R., Ometto, J. P. H. B., Pacheco, F. S., ... Marquina, S. (2015). Nitrogen management challenges in major watersheds of South America. *Environmental Research Letters*, 10, 065007. <https://doi.org/10.1088/1748-9326/10/6/065007>
- Campana, S. E. (1999). Chemistry and composition of fish otoliths: Pathways, mechanisms and applications. *Marine Ecology Progress Series*, 188, 263–297. <https://doi.org/10.3354/meps188263>
- Campana, S. E., & Neilson, J. D. (1985). Microstructure of fish otoliths. *Canadian Journal of Fisheries and Aquatic Sciences*, 42, 1014–1032. <https://doi.org/10.1139/f85-127>
- Chang, M.-Y., & Geffen, A. J. (2013). Taxonomic and geographic influences on fish otolith microchemistry. *Fish and Fisheries*, 14, 458–492. <https://doi.org/10.1111/j.1467-2979.2012.00482.x>
- Cheung, W. W. L., Brodeur, R. D., Okey, T. A., & Pauly, D. (2015). Projecting future changes in distributions of pelagic fish species of Northeast Pacific shelf seas. *Progress in Oceanography*, 130, 19–31. <https://doi.org/10.1016/j.pocean.2014.09.003>
- Dahlgren, C. P., Kellison, G. T., Adams, A. J., Gillanders, B. M., Kendall, M. S., Layman, C. A., ... Serafy, J. E. (2006). Marine nurseries and effective juvenile habitats: Concepts and applications. *Marine Ecology Progress Series*, 312, 291–295. <https://doi.org/10.3354/meps312291>
- de Matos, I. P., & Lucena, F. (2017). Descrição da pesca da pescada-amarela, *Cynoscion acoupa*, da costa do Pará. *Arquivos de Ciências do Mar*, 39, 66–73.
- Dingle, A., & Drake, V. A. (2007). What is migration? *BioScience*, 57, 113–121. <https://doi.org/10.1641/B570206>
- Dittmar, T., Hertkorn, N., Kattner, G., & Lara, R. J. (2006). Mangroves, a major source of dissolved organic carbon to the oceans. *Global Biogeochemical Cycles*, 20, GB1012
- Dittmar, T., & Lara, R. J. (2001). Driving forces behind nutrient and organic matter dynamics in a mangrove tidal creek in north Brazil. *Estuarine, Coastal and Shelf Science*, 52, 249–259. <https://doi.org/10.1006/ecss.2000.0743>
- Docherty, G., Jones, V., & Evershed, R. P. (2001). Practical and theoretical considerations in the gas chromatography/combustion/isotope ratio mass spectrometry $\delta^{13}\text{C}$ analysis of small polyfunctional compounds. *Rapid Communications in Mass Spectrometry*, 15, 730–738. [https://doi.org/10.1002/\(ISSN\)1097-0231](https://doi.org/10.1002/(ISSN)1097-0231)
- Elsdon, T. S., Ayvazian, S., McMahon, K. W., & Thorrold, S. R. (2010). Experimental evaluation of stable isotope fractionation in fish muscle and otoliths. *Marine Ecology Progress Series*, 408, 195–205. <https://doi.org/10.3354/meps08518>
- Elsdon, T. S., Wells, B. K., Campana, S. E., Gillanders, B. M., Jones, C. M., Limburg, K. E., ... Walther, B. D., (2008). Otolith chemistry to describe movements and life-history parameters of fishes: Hypotheses, assumptions, limitations and inferences. In R. N. Gibson, R. J. A. Atkinson, & J. D. M. Gordon (Eds.), *Oceanography and marine biology an annual review* (pp. 297–330). New York, NY: CRC Press.
- Ferreira, G. V. B., Barletta, M., Lima, A. R. A., Dantas, D. V., Justino, A. K. S., & Costa, M. F. (2016). Plastic debris contamination in the life cycle of *Acoupa* weakfish (*Cynoscion acoupa*) in a tropical estuary. *ICES Journal of Marine Science: Journal du Conseil*, 73, 2695–2707. <https://doi.org/10.1093/icesjms/fsw108>
- Geffen, A. J. (2012). Otolith oxygen and carbon stable isotopes in wild and laboratory-reared plaice (*Pleuronectes platessa*). *Environmental Biology of Fishes*, 95, 419–430. <https://doi.org/10.1007/s10641-012-0033-2>
- Gillanders, B. M., Able, K. W., Brown, J. A., Eggleston, D. B., & Sheridan, P. F. (2003). Evidence of connectivity between juvenile and adult habitats for mobile marine fauna: An important component of nurseries. *Marine Ecology Progress Series*, 247, 281–295. <https://doi.org/10.3354/meps247281>
- Grønkvær, P. (2016). Otoliths as individual indicators: A reappraisal of the link between fish physiology and otolith characteristics. *Marine and Freshwater Research*, 67, 881–888. <https://doi.org/10.1071/MF15155>
- Grønkvær, P., Pedersen, J. B., Ankjær, T. T., Kjeldsen, H., Heinemeier, J., Steingrund, P., ... Gillanders, B. M. (2013). Stable N and C isotopes in the organic matrix of fish otoliths: Validation of a new approach for studying spatial and temporal changes in the trophic structure of aquatic ecosystems. *Canadian Journal of Fisheries and Aquatic Sciences*, 70, 143–146. <https://doi.org/10.1139/cjfas-2012-0386>
- Halpern, B. S., Walbridge, S., Selkoe, K. A., Kappel, C. V., Micheli, F., D'Agrosa, C., ... Watson, R. (2008). A global map of human impact on marine ecosystems. *Science*, 319, 948–952. <https://doi.org/10.1126/science.1149345>
- Howland, M. R., Corr, L. T., Young, S. M. M., Jones, V., Jim, S., Van Der Merwe, N. J., ... Evershed, R. P. (2003). Expression of the dietary isotope signal in the compound-specific $\delta^{13}\text{C}$ values of pig bone lipids and amino acids. *International Journal of Osteoarchaeology*, 13, 54–65. [https://doi.org/10.1002/\(ISSN\)1099-1212](https://doi.org/10.1002/(ISSN)1099-1212)
- Huijbers, C. M., Nagelkerken, I., Debrot, A. O., & Jongejans, E. (2013). Geographic coupling of juvenile and adult habitat shapes spatial population dynamics of a coral reef fish. *Ecology*, 94, 1859–1870. <https://doi.org/10.1890/11-1759.1>
- Iglu, M. M., Nagelkerken, I., van der Velde, G., & Mgaya, Y. D. (2013). Mangrove fish production is largely fuelled by external food sources: A stable isotope analysis of fishes at the individual, species, and community levels from across the globe. *Ecosystems*, 16, 1336–1352. <https://doi.org/10.1007/s10021-013-9687-7>
- Jackson, J. B. C. (2001). Historical overfishing and the recent collapse of coastal ecosystems. *Science*, 293, 629–637. <https://doi.org/10.1126/science.1059199>
- Jamieson, R. E., Schwarcz, H. P., & Bratley, J. (2004). Carbon isotopic records from the otoliths of Atlantic cod (*Gadus morhua*) from eastern Newfoundland, Canada. *Fisheries Research*, 68, 83–97. <https://doi.org/10.1016/j.fishres.2004.02.009>
- Kalish, J. M. (1991). Oxygen and carbon stable isotopes in the otoliths of wild and laboratory-reared Australian salmon (*Arripis trutta*). *Marine Biology*, 110, 37–47. <https://doi.org/10.1007/BF01313090>
- Kimirei, I. A., Nagelkerken, I., Griffioen, B., Wagner, C., & Mgaya, Y. D. (2011). Ontogenetic habitat use by mangrove/seagrass-associated coral reef fishes shows flexibility in time and space. *Estuarine, Coastal and Shelf Science*, 92, 47–58. <https://doi.org/10.1016/j.ecss.2010.12.016>
- Kimirei, I. A., Nagelkerken, I., Slooter, N., Gonzalez, E. T., Huijbers, C. M., Mgaya, Y. D., & Rypel, A. L. (2015). Demography of fish populations reveals new challenges in appraising juvenile habitat values. *Marine Ecology Progress Series*, 518, 225–237. <https://doi.org/10.3354/meps11059>
- Knoppers, B., Ekau, W., & Figueiredo, A. G. (1999). The coast and shelf of east and northeast Brazil and material transport. *Geo-Marine Letters*, 19, 171–178. <https://doi.org/10.1007/s003670050106>
- Kruitwagen, G., Nagelkerken, I., Lugendo, B. R., Mgaya, Y. D., & Bonga, S. E. W. (2010). Importance of different carbon sources for macroinvertebrates and fishes of an interlinked mangrove–mudflat ecosystem (Tanzania). *Estuarine, Coastal and Shelf Science*, 88, 464–472. <https://doi.org/10.1016/j.ecss.2010.05.002>
- Krumme, U., Calderón, M. A., & Echterhoff, A. (2014). Intertidal migration of the four-eyed fish *Anableps anableps* in north Brazilian mangrove creeks. *Marine Ecology Progress Series*, 509, 271–287. <https://doi.org/10.3354/meps10863>
- Larsen, T., Taylor, D. L., Leigh, M. B., & O'Brien, D. M. (2009). Stable isotope fingerprinting: A novel method for identifying plant, fungal, or bacterial origins of amino acids. *Ecology*, 90, 3526–3535. <https://doi.org/10.1890/08-1695.1>
- Larsen, T., Ventura, M., Andersen, N., O'Brien, D. M., Piatkowski, U., & McCarthy, M. D. (2013). Tracing carbon sources through aquatic and terrestrial food webs using amino acid stable isotope fingerprinting. *PLoS ONE*, 8, e73441. <https://doi.org/10.1371/journal.pone.0073441>
- Larsen, T., Bach, L. T., Salvatelli, R., Wang, Y. V., Andersen, N., Ventura, M., & McCarthy, M. D. (2015). Assessing the potential of amino acid ^{13}C patterns as a carbon source tracer in marine sediments: Effects of algal growth conditions and sedimentary diagenesis. *Biogeosciences*, 12, 4979–4992. <https://doi.org/10.5194/bg-12-4979-2015>

- Lima, A. R. A., Barletta, M., & Costa, M. F. (2015). Seasonal distribution and interactions between plankton and microplastics in a tropical estuary. *Estuarine, Coastal and Shelf Science*, 165, 213–225. <https://doi.org/10.1016/j.ecss.2015.05.018>
- Loick-Wilde, N., Weber, S. C., Conroy, B. J., Capone, D. G., Coles, V. J., Medeiros, P. M., ... Montoya, J. P. (2016). Nitrogen sources and net growth efficiency of zooplankton in three Amazon River plume food webs. *Limnology and Oceanography*, 61, 460–481. <https://doi.org/10.1002/lno.10227>
- Lynam, C. P., Llope, M., Möllmann, C., Helaouët, P., Bayliss-Brown, G. A., & Stenseth, N. C. (2017). Interaction between top-down and bottom-up control in marine food webs. *Proceedings of the National Academy of Sciences of the United States of America*, 114, 1952–1957. <https://doi.org/10.1073/pnas.1621037114>
- McMahon, K. W., Berumen, M. L., Mateo, I., Elsdon, T. S., & Thorrold, S. R. (2011). Carbon isotopes in otolith amino acids identify residency of juvenile snapper (Family: Lutjanidae) in coastal nurseries. *Coral Reefs*, 30, 1135–1145. <https://doi.org/10.1007/s00338-011-0816-5>
- McMahon, K. W., Berumen, M. L., & Thorrold, S. R. (2012). Linking habitat mosaics and connectivity in a coral reef seascape. *Proceedings of the National Academy of Sciences*, 109, 15372–15376. <https://doi.org/10.1073/pnas.1206378109>
- McMahon, K. W., Fogel, M. L., Johnson, B. J., Houghton, L. A., Thorrold, S. R., & Gillanders, B. M. (2011). A new method to reconstruct fish diet and movement patterns from $\delta^{13}\text{C}$ values in otolith amino acids. *Canadian Journal of Fisheries and Aquatic Sciences*, 68, 1330–1340. <https://doi.org/10.1139/f2011-070>
- Melville, A. J., & Connolly, R. M. (2003). Spatial analysis of stable isotope data to determine primary sources of nutrition for fish. *Oecologia*, 136, 499–507. <https://doi.org/10.1007/s00442-003-1302-8>
- Nagelkerken, I., Russell, B. D., Gillanders, B. M., & Connell, S. D. (2015). Ocean acidification alters fish populations indirectly through habitat modification. *Nature Climate Change*, 6, 89–93.
- Nakazawa, T., & Yamamura, N. (2006). Breeding migration and population stability. *Population Ecology*, 49, 101–113.
- Nelson, J., Hanson, C. W., Koenig, C., & Chanton, J. (2011). Influence of diet on stable carbon isotope composition in otoliths of juvenile red drum *Sciaenops ocellatus*. *Aquatic Biology*, 13, 89–95. <https://doi.org/10.3354/ab00354>
- Nordhaus, I., Wolff, M., & Diele, K. (2006). Litter processing and population food intake of the mangrove crab *Ucides cordatus* in a high intertidal forest in northern Brazil. *Estuarine, Coastal and Shelf Science*, 67, 239–250. <https://doi.org/10.1016/j.ecss.2005.11.022>
- Pascal, P.-Y., Dubois, S. F., Goffette, A., & Lepoint, G. (2017). Influences of geothermal sulfur bacteria on a tropical coastal food web. *Marine Ecology Progress Series*, 578, 73–85. <https://doi.org/10.3354/meps12237>
- Pauly, D., & Christensen, V. (1995). Primary production required to sustain global fisheries. *Nature*, 374, 255–257. <https://doi.org/10.1038/374255a0>
- Radtke, R. L., Williams, D. F., & Hurley, P. C. F. (1987). The stable isotopic composition of bluefin tuna (*Thunnus thynnus*) otoliths: Evidence for physiological regulation. *Comparative Biochemistry and Physiology*, 87A, 797–801. [https://doi.org/10.1016/0300-9629\(87\)90403-8](https://doi.org/10.1016/0300-9629(87)90403-8)
- R Core Team (2017). R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. URL <https://www.R-project.org/>
- Santos, M. L. S., Medeiros, C., Muniz, K., Feitosa, F. A. N., Schwamborn, R., & Macêdo, S. J. (2008). Influence of the Amazon and Pará rivers on water composition and phytoplankton biomass on the adjacent shelf. *Journal of Coastal Research*, 243, 585–593. <https://doi.org/10.2112/05-0538.1>
- Schaeffer-Novelli, Y., Cintrón-Molero, G., Adame, R. R., de Camargo, T. M., Cintrón-Molero, G., & de Camargo, T. M. (1990). Variability of mangrove ecosystems along the Brazilian coast. *Estuaries*, 13, 204–218. <https://doi.org/10.2307/1351590>
- Schories, D., Barletta Bergan, A., Barletta, M., Krumme, U., Mehlig, U., & Rademaker, V. (2003). The keystone role of leaf-removing crabs in mangrove forests of North Brazil. *Wetlands Ecology and Management*, 11, 243–255. <https://doi.org/10.1023/A:1025011431984>
- Schwarcz, H. P., Gao, Y., Campana, S., Browne, D., Knyf, M., & Brand, U. (1998). Stable carbon isotope variations in otoliths of Atlantic cod (*Gadus morhua*). *Canadian Journal of Fisheries and Aquatic Sciences*, 55, 1798–1806. <https://doi.org/10.1139/f98-053>
- Secor, D., Henderson-Arzapalo, A., & Piccoli, P. M. (1995). Can otolith microchemistry chart patterns of migration and habitat utilization in anadromous fish? *Journal of Experimental Marine Biology and Ecology*, 192, 15–33. [https://doi.org/10.1016/0022-0981\(95\)00054-U](https://doi.org/10.1016/0022-0981(95)00054-U)
- Seitz, R. D., Wennhage, H., Bergstrom, U., Lipcius, R. N., & Ysebaert, T. (2014). Ecological value of coastal habitats for commercially and ecologically important species. *ICES Journal of Marine Science: Journal du Conseil*, 71, 648–665. <https://doi.org/10.1093/icesjms/fst152>
- Sheaves, M., Baker, R., Nagelkerken, I., & Connolly, R. M. (2015). True value of estuarine and coastal nurseries for fish: Incorporating complexity and dynamics. *Estuaries and Coasts*, 38, 401–414. <https://doi.org/10.1007/s12237-014-9846-x>
- Smith, W. O. Jr, & Demaster, D. J. (1996). Phytoplankton biomass and productivity in the Amazon River plume: Correlation with seasonal river discharge. *Continental Shelf Research*, 16, 291–319. [https://doi.org/10.1016/0278-4343\(95\)00007-N](https://doi.org/10.1016/0278-4343(95)00007-N)
- Solomon, C. T., Weber, P. K., Cech, J. J. Jr, Ingram, B. L., Conrad, M. E., Machavaram, M. V., ... Franklin, R. L. (2006). Experimental determination of the sources of otolith carbon and associated isotopic fractionation. *Canadian Journal of Fisheries and Aquatic Sciences*, 63, 79–89. <https://doi.org/10.1139/f05-200>
- Steffan, S. A., Chikaraishi, Y., Dharampal, P. S., Pauli, J. N., Guédot, C., & Ohkouchi, N. (2017). Unpacking brown food-webs: Animal trophic identity reflects rampant microbivory. *Ecology and Evolution*, 7, 3532–3541. <https://doi.org/10.1002/ece3.2951>
- Sturrock, A. M., Trueman, C. N., Milton, J. A., Waring, C. P., Cooper, M. J., & Hunter, E. (2014). Physiological influences can outweigh environmental signals in otolith microchemistry research. *Marine Ecology Progress Series*, 500, 245–264. <https://doi.org/10.3354/meps10699>
- Venables, W. N., & Ripley, B. D. (2002). *Modern applied statistics with S*. New York, NY: Springer. <https://doi.org/10.1007/978-0-387-21706-2>
- Vokhshoori, N. L., Larsen, T., & McCarthy, M. D. (2014). Reconstructing $\delta^{13}\text{C}$ isoscapes of phytoplankton production in a coastal upwelling system with amino acid isotope values of littoral mussels. *Marine Ecology Progress Series*, 504, 59–72. <https://doi.org/10.3354/meps10746>
- Walsh, R. G., He, S., & Yarnes, C. T. (2014). Compound-specific $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis of amino acids: A rapid, chloroformate-based method for ecological studies. *Rapid Communications in Mass Spectrometry*, 28, 96–108. <https://doi.org/10.1002/rcm.6761>
- Weber, S. C., Carpenter, E. J., Coles, V. J., Yager, P. L., Goes, J., & Montoya, J. P. (2017). Amazon River influence on nitrogen fixation and export production in the western tropical North Atlantic. *Limnology and Oceanography*, 62, 618–631. <https://doi.org/10.1002/lno.10448>
- Yi, X., Huang, Y., Zhuang, Y., Chen, H., Yang, F., Wang, W., ... Zhang, H. (2017). In situ diet of the copepod *Calanus sinicus* in coastal waters of the South Yellow Sea and the Bohai Sea. *Acta Oceanologica Sinica*, 36, 68–79. <https://doi.org/10.1007/s13131-017-0974-6>

How to cite this article: Vane K, Larsen T, Scholz-Böttcher BM, Kopke B, Ekau W. Ontogenetic resource utilization and migration reconstruction with $\delta^{13}\text{C}$ values of essential amino acids in the *Cynoscion acoupa* otolith. *Ecol Evol*. 2018;00:1–11. <https://doi.org/10.1002/ece3.4471>